

# ON HARDY SPACES OF (NONLOCAL) OPERATORS

KRZYSZTOF BOGDAN, BARTŁOMIEJ DYDA, AND TOMASZ LUKS

ABSTRACT. We characterize Hardy spaces and conditional Hardy spaces of the Laplacian and the fractional Laplacian by using Hardy-Stein type identities.

## 1. INTRODUCTION

Let  $D \subset \mathbb{R}^d$  be open and let  $x_0 \in D$ . For  $p > 0$  and  $0 < \alpha < 2$  we will consider the Hardy space  $\mathcal{H}^p(D, \alpha)$  of the fractional Laplacian  $\Delta^{\alpha/2}$ ,

$$(1) \quad \Delta^{\alpha/2}u(x) = \lim_{\eta \rightarrow 0^+} \int_{|y-x|>\eta} \mathcal{A} \frac{u(y) - u(x)}{|y-x|^{d+\alpha}} dy.$$

Here  $\mathcal{A} = \Gamma((d + \alpha)/2)/(2^{-\alpha}\pi^{d/2}|\Gamma(-\alpha/2)|)$ .  $\mathcal{H}^p(D, \alpha)$  is defined as follows. Let  $X$  be the isotropic  $\alpha$ -stable Lévy process ([9]), i.e. the symmetric Lévy process on  $\mathbb{R}^d$  with the Lévy measure  $\nu(dy) = \mathcal{A}|y|^{-d-\alpha}dy$  and zero Gaussian part ([9]). Let  $\mathbb{E}_x$  be the expectation for  $X$  starting at  $x \in \mathbb{R}^d$ . We define  $\tau_D = \inf\{t \geq 0 : X_t \notin D\}$ , the first exit time of  $X$  from  $D$ . A Borel function  $u : \mathbb{R}^d \rightarrow \mathbb{R}$  is called  $\alpha$ -harmonic on  $D$  if for every open  $U$  relatively compact in  $D$  (denoted  $U \subset\subset D$ ) we have

$$(2) \quad u(x) = \mathbb{E}_x u(X_{\tau_U}), \quad x \in U.$$

Here we assume that the expectation is absolutely convergent, in particular-finite. Equivalently,  $u$  is  $\alpha$ -harmonic on  $D$  if  $u$  is twice continuously differentiable on  $D$ ,  $\int_{\mathbb{R}^d} |u(y)|(1 + |y|)^{-d-\alpha}dy < \infty$ , and

$$(3) \quad \Delta^{\alpha/2}u(x) = 0, \quad x \in D.$$

---

*Date:* September 1, 2011.

*2010 Mathematics Subject Classification.* Primary 60J75; Secondary 60J50, 42B30, 31B25.

*Key words and phrases.* fractional Laplacian, Hardy spaces, conditional stable Lévy process.

First and second named authors were partially supported by MNiSW grant N N201 397137. Second named author was partially supported by the DFG through SFB-701 'Spectral Structures and Topological Methods in Mathematics'. Third named author was partially supported by Agence Nationale de la Recherche grant ANR-09-BLAN-0084-01 and by MNiSW grant N N201 373136.

We refer to [8] and [12] for this characterization and a detailed discussion of  $\alpha$ -harmonic functions, including structure theorems for nonnegative  $\alpha$ -harmonic functions, and explicit formulas for the Green function, Poisson kernel and Martin kernel of  $\Delta^{\alpha/2}$  for the ball. We also refer to remarks in [13, p. 120] explaining why the mean value property (2) is preferred over analogues of (3) for harmonic functions of more general Markov processes. The reader may verify, using (2) and the strong Markov property of  $X$ , that  $\{u(X_{\tau_U})\}_{U \subset \subset D}$  is a martingale ordered by inclusion of sets  $U$ . In particular,  $\mathbb{E}_x |u(X_{\tau_U})|^p$  is non-decreasing in  $U$ , if  $p \geq 1$ .

**Definition 1.** Let  $0 < p < \infty$ . We write  $u \in \mathcal{H}^p = \mathcal{H}^p(D, \alpha)$ , if  $u$  is  $\alpha$ -harmonic on  $D$  and

$$(4) \quad \|u\|_p := \sup_{U \subset \subset D} (\mathbb{E}_{x_0} |u(X_{\tau_U})|^p)^{1/p} < \infty.$$

The finiteness condition does not depend on the choice of  $x_0$ , because  $U \ni x \mapsto \mathbb{E}_x |u(X_{\tau_U})|^p$  satisfies Harnack inequality for arbitrary (Borel) function  $u$  ([12]). If  $p \leq q$ , then  $\mathcal{H}^p \supset \mathcal{H}^q$ .

We will say that nonnegative functions  $f(u)$  and  $g(u)$  are *comparable*, and write  $f(u) \asymp g(u)$ , if numbers  $0 < c \leq C < \infty$  exist such that  $cf(u) \leq g(u) \leq Cf(u)$  for every  $u$ .

The reader may notice that (4) is far from being explicit because it involves the distribution of  $X_{\tau_U}$  for all  $U \subset D$ . The following result and the exact formula for  $\|u\|_p$  given in (18) below may simplify the perspective. Let  $G_D(x, y)$  be the Green function of  $\Delta^{\alpha/2}$  for the Dirichlet problem on  $D$  ([12]).

**Theorem 1.** If  $1 < p < \infty$ , then  $\|u\|_p^p$  is comparable on  $\mathcal{H}^p$  with

$$(5) \quad |u(x_0)|^p + \int_D G_D(x_0, y) \int_{\mathbb{R}^d} \frac{[u(z) - u(y)]^2 (|u(z)| \vee |u(y)|)^{p-2}}{|z - y|^{d+\alpha}} dz dy.$$

In fact,  $u \in \mathcal{H}^p$  if and only if  $u$  is  $\alpha$ -harmonic in  $D$  and the integral is finite.

We will also describe Hardy spaces  $\mathcal{H}_h^p = \mathcal{H}_h^p(D, \alpha)$  related to the operator

$$(6) \quad \Delta_h^{\alpha/2}(u) = \frac{1}{h} \Delta^{\alpha/2}(hu).$$

Here  $h$  is a fixed function harmonic for  $\Delta^{\alpha/2}$ , positive on  $D$  and vanishing on  $D^c$ . The boundary behavior of functions in  $\mathcal{H}_h^p$  is of considerable interest because it directly relates to *ratios* of  $\alpha$ -harmonic functions. In fact, the above  $h$ -transform (6) of  $\Delta^{\alpha/2}$  is implicit in the relative Fatou theorems studied in [29, 24, 10] and in the theory of conditional stable processes [8, 14].

The paper is composed as follows. In Section 2 we observe the formula

$$(7) \quad \sup_{U \subset \subset D} \mathbb{E}_{x_0} u^2(X_{\tau_U}) = |u(x_0)|^2 + \int_D G_D(x_0, y) \int_{\mathbb{R}^d} \mathcal{A} \frac{[u(z) - u(y)]^2}{|z - y|^{d+\alpha}} dz dy$$

for the norm of  $\mathcal{H}^2$ , and we extend it to  $\mathcal{H}^p$  for  $p > 1$  in Lemma 4 and Theorem 1. The conditional Hardy spaces  $\mathcal{H}_h^p$  are characterized in Lemma 8, Theorem 2 and formula (35) in Section 3, see also Remark 1. In Section 4 we state analogous results for the classical Laplacian: formula (36) and Theorem 3. Formula (36) is the celebrated Hardy-Stein identity but Theorem 3 is apparently new, and may be considered a *conditional* Hardy-Stein identity. In Section 5 we describe the norm of the Hardy spaces in terms of Krickeberg decomposition for  $p \geq 1$ , and we prove a Littlewood-Paley inequality.

Formula (7) and its modifications (18, 35, 37) below are the main subject of the paper, and they may be considered nonlocal or conditional extensions of the classical Hardy-Stein equality, for which we refer the reader to (36) in Section 4 and [39, 32, 33].

Our work was motivated by the notion of the quadratic variation of martingales, carré du champ, and the characterization of the classical and martingale Hardy and Bergman spaces ([17, 28, 34, 33, 41, 39, 27, 42]). The resulting technique should apply to Hardy spaces of operators and Markov processes more general than the fractional Laplacian and the isotropic stable Lévy process. The inclusion of both jump and continuous processes in the present paper is intended to clarify the methodology and indicate such extensions. Our development is analytic, except for the *definitions* of the Hardy spaces, which are probabilistic. These too, however, can be formulated analytically by using the harmonic measures of the Laplacian and the fractional Laplacian ([4, 26]).

## 2. CHARACTERIZATION OF $\mathcal{H}^p$

Consider open  $U \subset\subset D$  and a real-valued function  $\phi : \mathbb{R}^d \rightarrow \mathbb{R}$  which is  $C^2$  in a neighborhood of  $\bar{U}$  and satisfies  $\int_{\mathbb{R}^d} |\phi(y)|(1 + |y|)^{-d-\alpha} < \infty$ . Then  $\Delta^{\alpha/2}\phi$  is bounded on  $\bar{U}$ , and for every  $x \in \mathbb{R}^d$  we have

$$(8) \quad \phi(x) = \mathbb{E}_x \phi(X_{\tau_U}) - \int_U G_U(x, y) \Delta^{\alpha/2} \phi(y) dy.$$

Indeed, the difference of both sides of (8) vanishes continuously on  $U^c$  and annihilates  $\Delta^{\alpha/2}$  on  $U$ . Thus the difference is zero on  $U$  by the maximum principle for  $\Delta^{\alpha/2}$ , see [8, Lemma 3.8 and the proof of Theorem 3.9], or [11, Appendix].

**Lemma 1.** *If  $u$  is  $\alpha$ -harmonic on  $D$  and  $U \subset\subset D$ , then*

$$(9) \quad \mathbb{E}_{x_0} u^2(X_{\tau_U}) = u(x_0)^2 + \int_U G_U(x_0, y) \int_{\mathbb{R}^d} \mathcal{A} \frac{[u(z) - u(y)]^2}{|z - y|^{d+\alpha}} dz dy.$$

*Proof.* If  $\int_{\mathbb{R}^d} u(y)^2(1 + |y|)^{-d-\alpha} = \infty$ , then  $\int_{\mathbb{R}^d} [u(z) - u(y)]^2 / |z - y|^{d+\alpha} dz = \infty$  for every  $y$ . Also  $\mathbb{E}_{x_0} u^2(X_{\tau_U}) = \infty$ , because the distribution of  $X_{\tau_U}$  has density function bounded below by a multiple of  $(1 + |y|)^{-d-\alpha}$  in the complement of

the neighborhood of  $\bar{U}$  ([5]). Therefore in what follows we may assume that  $\int_{\mathbb{R}^d} u(y)^2(1 + |y|)^{-d-\alpha} < \infty$ . Since  $u^2$  is smooth in a neighborhood of  $\bar{U}$ ,  $\Delta^{\alpha/2}(u^2)$  is bounded on  $\bar{U}$ . By (8) with  $\phi = u^2$ , for  $x \in \mathbb{R}^d$  we have

$$(10) \quad \mathbb{E}_x u^2(X_{\tau_U}) = u^2(x) + \int_U G_U(x, y) \Delta^{\alpha/2}(u^2)(y) dy.$$

For  $y \in \bar{U}$ ,  $z \in \mathbb{R}^d$ , we have  $u^2(z) - u^2(y) - 2u(y)[u(z) - u(y)] = [u(z) - u(y)]^2$ . Since  $\Delta^{\alpha/2}u(y) = 0$ , we have

$$\begin{aligned} \Delta^{\alpha/2}u^2(y) &= \Delta^{\alpha/2}u^2(y) - 2u(y)\Delta^{\alpha/2}u(y) \\ &= \lim_{\eta \rightarrow 0^+} \int_{\mathcal{A}} \frac{u^2(z) - u^2(y) - 2u(y)[u(z) - u(y)]}{|z - y|^{d+\alpha}} dz \\ &= \int_{\mathbb{R}^d} \mathcal{A} \frac{[u(z) - u(y)]^2}{|z - y|^{d+\alpha}} dz, \end{aligned}$$

and (9) follows.  $\square$

Recall that  $G_U(x, y) \uparrow G_D(x, y)$  as  $U \uparrow D$ . By the monotone convergence theorem we obtain the description of  $\mathcal{H}^2$  aforementioned in Introduction.

**Corollary 2.** *If  $u$  is  $\alpha$ -harmonic in  $D$ , then (7) holds.*

We conclude that  $\mathcal{H}^2$  consists of precisely all those functions  $\alpha$ -harmonic on  $D$  for which the quadratic form on the right hand side of (7) is finite.

We will now consider (general)  $p > 1$ . We note that  $x \mapsto |x|^p$  is convex on  $\mathbb{R}$ , with the derivative  $pa|a|^{p-2}$  at  $x = a$ . For  $a, b \in \mathbb{C}$  we let

$$(11) \quad F(a, b) = |b|^p - |a|^p - p\bar{a}|a|^{p-2}(b - a),$$

a second order Taylor remainder. We have  $F(a, b) = |b|^p$  if  $a = 0$  and  $F(a, b) = (p - 1)|a|^p$  if  $b = 0$ . By convexity,  $F(a, b) \geq 0$  for  $a, b \in \mathbb{R}$ .

**Example 1.** For (even)  $p = 2, 4, \dots$  and  $a, b \in \mathbb{R}$ , we have

$$F(a, b) = b^p - a^p - pa^{p-1}(b - a) = (b - a)^2 \sum_{k=0}^{p-2} (k + 1)b^{p-2-k}a^k.$$

For  $p \in (1, 2)$  we will use a certain approximation procedure. Let  $\varepsilon, b$  and  $a$  be real numbers, and define

$$(12) \quad F_\varepsilon(a, b) = \operatorname{Re} F(a + i\varepsilon, b + i\varepsilon) = |b + i\varepsilon|^p - |a + i\varepsilon|^p - pa|a + i\varepsilon|^{p-2}(b - a).$$

Of course,  $F_\varepsilon(a, b) \rightarrow F(a, b)$  as  $\varepsilon \rightarrow 0$ .

**Lemma 3.** *For every  $p > 1$ , we have*

$$(13) \quad F(a, b) \asymp (b - a)^2(|b| \vee |a|)^{p-2}, \quad a, b \in \mathbb{R}.$$

*If  $p \in (1, 2)$ , then uniformly in  $\varepsilon, a, b \in \mathbb{R}$ , we have*

$$(14) \quad F_\varepsilon(a, b) \leq c(b - a)^2(|b| \vee |a|)^{p-2}.$$

*Proof.* We denote  $K(a, b) = (b - a)^2(|b| \vee |a|)^{p-2}$ . For every  $k \in \mathbb{R}$ ,  $F(ka, kb) = |k|^p F(a, b)$  and  $K(ka, kb) = |k|^p K(a, b)$ . If  $a = 0$ , then (13) becomes equality, hence we may assume that  $a \neq 0$ , in fact – that  $a = 1$ . Let  $f(b) = F(1, b) = |b|^p - 1 - p(b - 1)$ . We will prove that

$$(15) \quad c_p(b - 1)^2(|b| \vee 1)^{p-2} \leq f(b) \leq C_p(b - 1)^2(|b| \vee 1)^{p-2}.$$

Since  $f(1) = f'(1) = 0$  and  $f''(y) = p(p - 1)|y|^{p-2}$ , we obtain

$$(16) \quad f(b) = p(p - 1) \int_1^b \int_1^x |y|^{p-2} dy dx = p(p - 1) \int_1^b |y|^{p-2} (b - y) dy.$$

The first integral is over a simplex of area  $(b - 1)^2/2$ , and it is a monotone function of the simplex (as ordered by inclusion). For  $b$  close to 1 the integral is comparable to  $(b - 1)^2$ . For large  $|b|$  the (second) integral is comparable to  $|b|^p$ . This proves (15), hence (13). We will now prove (14) for  $p \in (1, 2)$ . Let  $f_\varepsilon(b) = F_\varepsilon(1, b) = (b^2 + \varepsilon)^{p/2} - (1 + \varepsilon^2)^{p/2} - p(1 + \varepsilon^2)^{(p-2)/2}(b - 1)$ . We have  $f_\varepsilon(1) = f'_\varepsilon(1) = 0$  and

$$f''_\varepsilon(b) = (b^2 + \varepsilon^2)^{(p-4)/2} p[b^2(p - 1) + \varepsilon^2] \asymp (b^2 + \varepsilon^2)^{(p-2)/2} \leq |b|^{p-2}.$$

Thus, uniformly in  $\varepsilon$ ,

$$f_\varepsilon(b) \leq c \int_1^b \int_1^x |y|^{p-2} dy dx = c \int_1^b (b - y)|y|^{p-2} dy \asymp (b - 1)^2(|b| \vee 1)^{p-2}.$$

If  $a \neq 0$ , then  $F_\varepsilon(a, b) = |a|^p f_{\varepsilon/a}(b/a) \leq c|a|^p (b/a - 1)^2(|b/a| \vee 1)^{p-2}$ . If  $a = 0$ , then  $F_\varepsilon(a, b) = (b^2 + \varepsilon^2)^{p/2} - |\varepsilon|^p \asymp |\varepsilon|^p [(b/\varepsilon)^2 \wedge |b/\varepsilon|^p] \leq |b|^p$ . The proof of (14) is complete.  $\square$

**Lemma 4.** *If  $u$  is  $\alpha$ -harmonic in  $D$ ,  $U \subset\subset D$ , and  $p > 1$ , then*

$$(17) \quad \mathbb{E}_{x_0} |u(X_{\tau_U})|^p = |u(x_0)|^p + \int_U G_U(x_0, y) \int_{\mathbb{R}^d} \mathcal{A} \frac{F(u(y), u(z))}{|z - y|^{d+\alpha}} dz dy.$$

*Proof.* We proceed as in Lemma 1. In particular we will assume that  $\int_{\mathbb{R}^d} |u(y)|^p (1 + |y|)^{-d-\alpha} < \infty$ , for otherwise both sides of (17) are infinite. We will first consider the case of  $p \geq 2$  and apply (8) to  $\phi = |u|^p \in C^2(D)$ . For  $y \in D$  we have

$\Delta^{\alpha/2}u(y) = 0$ , hence

$$\begin{aligned} \Delta^{\alpha/2}|u|^p(y) &= \Delta^{\alpha/2}|u|^p(y) - pu(y)|u(y)|^{p-2}\Delta^{\alpha/2}u(y) \\ &= \lim_{\eta \rightarrow 0^+} \int_{\{z \in \mathbb{R}^d: |z-y| > \eta\}} \mathcal{A} \frac{|u(z)|^p - |u(y)|^p - pu(y)|u(y)|^{p-2}[u(z) - u(y)]}{|z-y|^{d+\alpha}} dz \\ &= \int_{\mathbb{R}^d} \mathcal{A} \frac{F(u(y), u(z))}{|z-y|^{d+\alpha}} dz. \end{aligned}$$

This and (8) yield (17) for  $p \geq 2$ . We now consider  $1 < p < 2$ . We note that  $|u + i\varepsilon|^p \in C^2(D)$ . As in the first part of the proof,

$$\begin{aligned} \Delta^{\alpha/2}|u + i\varepsilon|^p(y) &= \Delta^{\alpha/2}|u + i\varepsilon|^p(y) - pu(y)|u(y) + i\varepsilon|^{p-2}\Delta^{\alpha/2}u(y) \\ &= \int_{\mathbb{R}^d} \mathcal{A} \frac{F_\varepsilon(u(y), u(z))}{|z-y|^{d+\alpha}} dz, \end{aligned}$$

hence

$$\mathbb{E}_{x_0}|u(X_{\tau_U}) + i\varepsilon|^p = |u(x_0) + i\varepsilon|^p + \int_U G_U(x_0, y) \int_{\mathbb{R}^d} \mathcal{A} \frac{F_\varepsilon(u(y), u(z))}{|z-y|^{d+\alpha}} dz dy.$$

We let  $\varepsilon \rightarrow 0$ . By Fatou's lemma, (14), (13) and dominated convergence, we obtain (17).  $\square$

*Proof of Theorem 1.* Lemma 3, Lemma 4 and monotone convergence imply the comparability of  $\|u\|_p^p$  and (5) with the same constants as in (15), under the mere assumption that  $u$  be  $\alpha$ -harmonic on  $D$ . In fact,

$$(18) \quad \|u\|_p^p = |u(x_0)|^p + \int_D G_D(x_0, y) \int_{\mathbb{R}^d} \mathcal{A} \frac{F(u(y), u(z))}{|z-y|^{d+\alpha}} dz dy. \quad \square$$

We note that in many cases the exact asymptotics of  $G_D$  is known. For instance, if  $D$  is a bounded  $C^{1,1}$  domain in  $\mathbb{R}^d$  and  $d > \alpha$ , then

$$G_D(x_0, y) \asymp \delta_D(y)^{\alpha/2} |y - x_0|^{\alpha-d},$$

where  $\delta_D(y) := \text{dist}(y, D^c)$ . Here we refer the reader to [22], also for the case of Lipschitz domains.

Recall the definition of  $F_\varepsilon$ , (12), and the fact that  $F_0 = F$  of (11). Before moving to conditional Hardy spaces we will record the following observation.

**Lemma 5.** *For every  $p > 1$  and  $a_1, a_2, b_1, b_2, \varepsilon \in \mathbb{R}$ , we have*

$$(19) \quad F_\varepsilon(a_1 \wedge a_2, b_1 \wedge b_2) \leq F_\varepsilon(a_1, b_1) \vee F_\varepsilon(a_2, b_2),$$

$$(20) \quad F_\varepsilon(a_1 \vee a_2, b_1 \vee b_2) \leq F_\varepsilon(a_1, b_1) \vee F_\varepsilon(a_2, b_2).$$

*In particular,  $F(a \wedge 1, b \wedge 1) \leq F(a, b)$  and  $F(a \vee (-1), b \vee (-1)) \leq F(a, b)$ , for all  $a, b \in \mathbb{R}$ . The latter also extends to  $K(a, b) = (b - a)^2(|b| \vee |a|)^{p-2}$ .*

*Proof.* Let  $\varepsilon \neq 0$ . We claim that the function  $b \mapsto F_\varepsilon(a, b)$  decreases on  $(-\infty, a]$  and increases on  $[a, \infty)$ . To see this, we consider

$$(21) \quad \frac{\partial F_\varepsilon}{\partial b}(a, b) = pb(b^2 + \varepsilon^2)^{p/2-1} - pa(a^2 + \varepsilon^2)^{p/2-1}.$$

The function  $h(x) = px(x^2 + \varepsilon^2)^{p/2-1}$  has derivative  $h'(x) = p(x^2 + \varepsilon^2)^{p/2-2}(x^2(p-1) + \varepsilon^2) > 0$ . It follows that the difference in (21) is positive if  $b > a$  and negative if  $b < a$ . This proves our claim.

Furthermore the function  $a \mapsto F_\varepsilon(a, b)$  decreases on  $(-\infty, b]$  and increases on  $[b, \infty)$ , as follows from calculating the derivative,

$$(22) \quad \frac{\partial F_\varepsilon}{\partial a}(a, b) = p(a-b)(a^2 + \varepsilon^2)^{p/2-2}(a^2(p-1) + \varepsilon^2).$$

We will now prove (19). If  $b_1 \wedge b_2 = b_1$  and  $a_1 \wedge a_2 = a_1$  (or  $b_1 \wedge b_2 = b_2$  and  $a_1 \wedge a_2 = a_2$ ), then (19) is trivial. If  $b_1 \wedge b_2 = b_1$  and  $a_1 \wedge a_2 = a_2$ , then the monotonicity of  $F$  yields

$$\begin{aligned} F_\varepsilon(a_2, b_1) &\leq F_\varepsilon(a_1, b_1), & \text{if } b_1 < a_2, \\ F_\varepsilon(a_2, b_1) &\leq F_\varepsilon(a_2, b_2), & \text{if } b_1 \geq a_2. \end{aligned}$$

The case  $b_1 \wedge b_2 = b_2$  and  $a_1 \wedge a_2 = a_1$  obtains by renaming the arguments. This proves inequality (19). (20) follows from (19) and the identity

$$F_\varepsilon(-a, -b) = F_\varepsilon(a, b).$$

The case  $\varepsilon = 0$  obtains by passing to the limit. When  $a = b$ , we have  $F(a, b) = 0$ , which yields the second last statement of the lemma. For  $K$  we obviously have  $(b \wedge 1 - a \wedge 1)^2(|b \wedge 1| \vee |a \wedge 1|)^{p-2} \leq (b-a)^2(|b| \vee |a|)^{p-2}$  and  $(b \vee (-1) - a \vee (-1))^2(|b \vee (-1)| \vee |a \vee (-1)|)^{p-2} \leq (b-a)^2(|b| \vee |a|)^{p-2}$ .  $\square$

In passing we note that if the right-hand side of (17) is finite for  $u$ , then it is also finite (in fact smaller) for  $u \wedge 1$  and  $u \vee (-1)$ . The latter functions have smaller values and increments than  $u$ , a property defining *normal contractions* for Dirichlet forms ([19]).

### 3. CHARACTERIZATION OF $\mathcal{H}_h^2$

The fractional Laplacian is a nonlocal operator and the corresponding stochastic process  $X$  has jumps. In consequence the definitions of  $\alpha$ -harmonicity (2) and (3) involve the values of the function on the whole of  $D^c$  ([12]). This is somewhat unusual compared with the classical theory of the Laplacian and the Brownian motion, and attempts were made by various authors to ascribe genuine boundary conditions to such processes and functions ([3, 7, 21, 29]). One possibility is to study the boundary behavior of harmonic functions after an appropriate normalization. In this section we will focus on  $\alpha$ -harmonic functions vanishing on  $D^c$ , so that  $D^c$  may be ignored, and we will use Doob's

$h$ -conditioning to normalize ([16]). Let  $h$  be harmonic and positive on  $D$ , and let  $h$  vanish on  $D^c$ . In particular  $h$  may be a Martin integral,

$$h(x) = \int M(x, z) \mu(dz), \quad x \in D,$$

where  $\mu \geq 0$  is a finite nonzero Borel measure on the set of the so-called accessible points of  $\partial D$ , and  $M(\cdot, z)$  denotes the Martin kernel with the pole at  $z \in \partial D$ , see [12]. We like to note that if  $D$  is Lipschitz, then all the points of  $\partial D$  are accessible ([12]), while in the general case  $\partial_M D \subset \partial D \cup \{\infty\}$ . Doob's  $h$ -transform (6) yields a nontrivial modification of the potential theory of  $\Delta^{\alpha/2}$ . In particular, the ratio  $u/h$  is harmonic for  $\Delta_h^{\alpha/2}$  if  $u$  is  $\alpha$ -harmonic and vanishes on  $D^c$  (see below). This observation offers a convenient framework for studying growth and boundary behavior of such ratios.

We will consider the transition semigroup

$$(23) \quad P_t^h f(x) = \frac{1}{h(x)} \int p_D(t, x, y) f(y) h(y) dy.$$

Here  $p_D(t, x, y) = \mathbb{E}_x\{t < \tau_D : p(t - \tau_D, X_{\tau_D}, y)\}$  is the time-homogeneous transition density of  $X$  killed on leaving  $D$  ([8]). The semigroup property of  $P_t^h$  follows directly from the Chapman-Kolmogorov equations for  $p_D$ ,

$$\int_{\mathbb{R}^d} p_D(s, x, y) p_D(t, y, z) dy = p_D(s + t, x, z).$$

By  $\alpha$ -harmonicity and the optional stopping theorem,  $\mathbb{E}_x h(X_{\tau_U \wedge t}) = h(x)$ , if  $x \in U \subset\subset D$ . Letting  $U \uparrow D$ , by Fatou's lemma we obtain  $\int p_D(t, x, y) h(y) dy = \mathbb{E}_x\{t < \tau_D : h(X_t)\} \leq h(x)$ , i.e.  $P_t^h$  is subprobabilistic.

The conditional process is defined as the Markov process with the transition semigroup  $P^h$ , and it will be denoted by the same symbol  $X$ . We let  $\mathbb{E}_x^h$  be the corresponding expectation for  $X$  starting at  $x \in D$ ,

$$\mathbb{E}_x^h f(X_t) = \frac{1}{h(x)} \mathbb{E}_x[t < \tau_D; f(X_t) h(X_t)],$$

see also [8]. A Borel function  $r : D \rightarrow \mathbb{R}$  is  $h$ -harmonic (on  $D$ ) if and only if for every open  $U \subset\subset D$  we have

$$(24) \quad r(x) = \mathbb{E}_x^h r(X_{\tau_U}) = \frac{1}{h(x)} \mathbb{E}_x[X_{\tau_U} \in D; r(X_{\tau_U}) h(X_{\tau_U})], \quad x \in U.$$

Here we assume that the expectation is absolutely convergent, in particular—finite. It is evident that  $r$  is  $h$ -harmonic if and only if  $r = u/h$  on  $D$ , where  $u$  is *singular  $\alpha$ -harmonic on  $D$*  ([12]), i.e.  $u$  is  $\alpha$ -harmonic on  $D$  and  $u = 0$  on  $D^c$ . We are interested in  $L^p$  integrability of  $u/h$ , since it amounts to *weighted  $L^p$  integrability of  $u$* . The following definition is adapted from [30].

**Definition 2.** For  $0 < p < \infty$  we define  $\mathcal{H}_h^p = \mathcal{H}_h^p(D, \alpha)$  as the class of all the functions  $u : \mathbb{R}^d \rightarrow \mathbb{R}$ , singular  $\alpha$ -harmonic on  $D$ , and such that

$$\|u\|_{\mathcal{H}_h^p}^p := \sup_{U \subset \subset D} \mathbb{E}_{x_0}^h \left| \frac{u(X_{\tau_U})}{h(X_{\tau_U})} \right|^p = \frac{1}{h(x_0)} \sup_{U \subset \subset D} \mathbb{E}_{x_0} \frac{|u(X_{\tau_U})|^p}{h(X_{\tau_U})^{p-1}} < \infty,$$

where  $\mathbb{E}_{x_0} |u(X_{\tau_U})|^p / h(X_{\tau_U})^{p-1}$  means  $\mathbb{E}_{x_0} [X_{\tau_U} \in D; |u(X_{\tau_U})|^p / h(X_{\tau_U})^{p-1}]$ .

By Harnack inequality,  $\mathcal{H}_h^p$  does not depend on the choice of  $x_0 \in D$ .

**Remark 1.** Note that the elements of this  $\mathcal{H}_h^p$  are  $\alpha$ -harmonic, rather than  $h$ -harmonic. In view of Definition 2, the genuine conditional Hardy space of  $\Delta^{\alpha/2}$  is  $\frac{1}{h}\mathcal{H}_h^p = \{u/h : u \in \mathcal{H}_h^p\}$ , with the norm  $\|u/h\| = \|u\|_{\mathcal{H}_h^p}$ .  $\mathcal{H}_h^p$  is more of a weighted Hardy space of  $\Delta^{\alpha/2}$ , but it is convenient to call it the *conditional Hardy space*, too. Below we will focus on calculating  $\|u\|_{\mathcal{H}_h^p}$ , which yields a description of both spaces.

**Lemma 6.** If  $u$  is singular  $\alpha$ -harmonic on  $D$  and  $U \subset \subset D$ , then

$$(25) \quad \mathbb{E}_{x_0} \frac{u^2(X_{\tau_U})}{h(X_{\tau_U})} = \frac{u(x_0)^2}{h(x_0)} + \int_U G_U(x_0, y) \int_{\mathbb{R}^d} \mathcal{A} \left[ \frac{u(z)}{h(z)} - \frac{u(y)}{h(y)} \right]^2 \frac{h(z) dz dy}{|z - y|^{d+\alpha}}.$$

*Proof.* Let  $y \in D$ . For arbitrary  $z \in \mathbb{R}^d$  we have

$$(26) \quad \begin{aligned} & \frac{u^2(z)}{h(z)} - \frac{u^2(y)}{h(y)} - 2 \frac{u(y)}{h(y)} (u(z) - u(y)) + \frac{u^2(y)}{h^2(y)} (h(z) - h(y)) \\ &= \left[ \frac{u(z)}{h(z)} - \frac{u(y)}{h(y)} \right]^2 h(z). \end{aligned}$$

By (26) and  $\alpha$ -harmonicity,

$$(27) \quad \begin{aligned} \Delta^{\alpha/2} \left( \frac{u^2}{h} \right) (y) &= \Delta^{\alpha/2} \left( \frac{u^2}{h} \right) (y) - 2 \frac{u(y)}{h(y)} \Delta^{\alpha/2} u(y) + \frac{u^2(y)}{h^2(y)} \Delta^{\alpha/2} h(y) \\ &= \int_{\mathbb{R}^d} \mathcal{A} \left[ \frac{u(z)}{h(z)} - \frac{u(y)}{h(y)} \right]^2 |z - y|^{-d-\alpha} h(z) dz. \end{aligned}$$

Noteworthy, the integrand is nonnegative. Following (10), for  $u^2/h$  we get

$$\mathbb{E}_{x_0} \frac{u^2(X_{\tau_U})}{h(X_{\tau_U})} = \frac{u(x_0)^2}{h(x_0)} + \int_U G_U(x_0, y) \Delta^{\alpha/2} \left( \frac{u^2}{h} \right) (y).$$

By using (27) we obtain (25).  $\square$

We can interpret (25) in terms of  $h$ -conditioning and  $r = u/h$  as follows,

$$(28) \quad \mathbb{E}_{x_0}^h r(X_{\tau_U})^2 = r(x_0)^2 + \int_U G_U(x_0, y) \int_{\mathbb{R}^d} \mathcal{A} \frac{[r(z) - r(y)]^2}{|z - y|^{d+\alpha}} \frac{h(z)}{h(x_0)} dz dy.$$

This is an analogue of (7), and also indicates the general situation. For  $p > 1$  we will consider the expressions of the form

$$F\left(\frac{a}{s}, \frac{b}{t}\right), \quad a, b \in \mathbb{C}, \quad s, t > 0,$$

see (11). If  $1 < p < 2$ , then we will need an approximation based on

$$F_\varepsilon\left(\frac{a}{s}, \frac{b}{t}\right), \quad a, b \in \mathbb{C}, \quad s, t > 0, \quad \varepsilon \in \mathbb{R},$$

where  $F_\varepsilon$  is defined in (12). By Lemma 3 we have that  $F(a/s, b/t) \geq 0$ ,

$$(29) \quad F\left(\frac{a}{s}, \frac{b}{t}\right) \asymp \left(\frac{b}{t} - \frac{a}{s}\right)^2 \left(\frac{|b|}{t} \vee \frac{|a|}{s}\right)^{p-2}, \quad a, b \in \mathbb{R}, \quad s, t > 0,$$

and uniformly in  $\varepsilon \in \mathbb{R}$  we have

$$(30) \quad F_\varepsilon\left(\frac{a}{s}, \frac{b}{t}\right) \leq c \left(\frac{b}{t} - \frac{a}{s}\right)^2 \left(\frac{|b|}{t} \vee \frac{|a|}{s}\right)^{p-2}, \quad a, b \in \mathbb{R}, \quad s, t > 0,$$

and the comparisons in (29) hold with the constants  $c_p$  and  $C_p$  of (15).

**Lemma 7.** *For  $p > 1$ ,  $a, b \in \mathbb{C}$  and  $s, t > 0$ , we have*

$$(31) \quad F\left(\frac{a}{s}, \frac{b}{t}\right) = \frac{|b|^p}{t^p} - \frac{|a|^p}{ts^{p-1}} - \frac{p|a|^{p-2}\bar{a}(b-a)}{ts^{p-1}} + \frac{(p-1)|a|^p(t-s)}{ts^p}.$$

*Proof.* The equality is straightforward to verify, and is left to the reader.  $\square$

The homogeneity seen on the left-hand side of (31) is an interesting feature for the right-hand side of (31). We also like to note that  $tF(a/s, b/t)$  is the remainder in the first order Taylor's expansion of  $(a, s) \mapsto |a|^p/s^{p-1}$  at  $(b, t)$  and, of course,  $F_\varepsilon(a/s, b/t) \rightarrow F(a/s, b/t)$  as  $\varepsilon \rightarrow 0$ .

**Lemma 8.** *If  $u$  is singular  $\alpha$ -harmonic on  $D$ ,  $U \subset\subset D$  and  $p > 1$ , then*

$$\mathbb{E}_{x_0} \frac{|u(X_{\tau_U})|^p}{h(X_{\tau_U})^{p-1}} = \frac{|u(x_0)|^p}{h(x_0)^{p-1}} + \int_U G_U(x_0, y) \int_{\mathbb{R}^d} F\left(\frac{u(y)}{h(y)}, \frac{u(z)}{h(z)}\right) \mathcal{A} \frac{h(z)dzdy}{|z-y|^{d+\alpha}}.$$

*Proof.* If  $p \geq 2$ , then  $|u|^p/h^{p-1} \in C^2(D)$ . By (10),

$$(32) \quad \mathbb{E}_{x_0} \frac{|u(X_{\tau_U})|^p}{h(X_{\tau_U})^{p-1}} = \frac{|u(x_0)|^p}{h(x_0)^{p-1}} + \int_U G_U(x_0, y) \Delta^{\alpha/2} \left(\frac{|u|^p}{h^{p-1}}\right)(y) dy.$$

By  $\alpha$ -harmonicity of  $h$  and  $u$ , and by Lemma 7,

$$\begin{aligned}
\Delta^{\alpha/2} \left( \frac{|u|^p}{h^{p-1}} \right) (y) &= \Delta^{\alpha/2} \left( \frac{|u|^p}{h^{p-1}} \right) (y) - \frac{p|u(y)|^{p-2}u(y)}{h(y)^{p-1}} \Delta^{\alpha/2} u(y) \\
&\quad + \frac{(p-1)|u(y)|^p}{h(y)^p} \Delta^{\alpha/2} h(y) \\
&= \lim_{\eta \rightarrow 0^+} \int_{\{z \in \mathbb{R}^d: |z-y| > \eta\}} \mathcal{A} \left[ \frac{|u(z)|^p}{h(z)^{p-1}} - \frac{|u(y)|^p}{h(y)^{p-1}} - \frac{p|u(y)|^{p-2}u(y)}{h(y)^{p-1}} (u(z) - u(y)) \right. \\
&\quad \left. + \frac{(p-1)|u(y)|^p}{h(y)^p} (h(z) - h(y)) \right] |z-y|^{-d-\alpha} dz \\
(33) \quad &= \int_{\mathbb{R}^d} h(z) F(u(y)/h(y), u(z)/h(z)) \mathcal{A} |z-y|^{-d-\alpha} dz,
\end{aligned}$$

see (31). This gives the result for  $p \geq 2$ . If  $1 < p < 2$  then we argue as follows. Let  $\varepsilon \rightarrow 0$  and consider  $u + i\varepsilon h$  in place of  $u$  in (32). By  $\alpha$ -harmonicity,  $\Delta^{\alpha/2} (|u + i\varepsilon h|^p h^{1-p})(y)$  equals

$$\begin{aligned}
&\lim_{\eta \rightarrow 0^+} \int_{\{z \in \mathbb{R}^d: |z-y| > \eta\}} \mathcal{A} \left[ \frac{|u(z) + i\varepsilon h(z)|^p}{h(z)^{p-1}} - \frac{|u(y) + i\varepsilon h(y)|^p}{h(y)^{p-1}} \right. \\
&\quad \left. - \frac{p|u(y) + i\varepsilon h(y)|^{p-2}u(y)}{h(y)^{p-1}} (u(z) - u(y)) \right. \\
&\quad \left. + \frac{(p-1)|u(y) + i\varepsilon h(y)|^p}{h(y)^p} (h(z) - h(y)) \right] |z-y|^{-d-\alpha} dz \\
(34) \quad &= \int_{\mathbb{R}^d} h(z) F_\varepsilon(u(y)/h(y), u(z)/h(z)) \mathcal{A} |z-y|^{-d-\alpha} dz.
\end{aligned}$$

We then use Lemma 7, Fatou's lemma, and dominated convergence.  $\square$

**Theorem 2.** *Let  $1 < p < \infty$ . For  $u$  singular  $\alpha$ -harmonic on  $D$ ,  $\|u\|_{\mathcal{H}_h^p}^p$  is comparable with*

$$\frac{|u(x_0)|^p}{h(x_0)^p} + \int_D \frac{G_D(x_0, y)}{h(x_0)} \int_{\mathbb{R}^d} \left( \frac{|u(z)|}{h(z)} \vee \frac{|u(y)|}{h(y)} \right)^{p-2} \left[ \frac{u(z)}{h(z)} - \frac{u(y)}{h(y)} \right]^2 \frac{h(z) dz dy}{|z-y|^{d+\alpha}}.$$

*Proof.* The result follows from Lemma 8 and Lemma 7. In fact,

$$(35) \quad \|u\|_{\mathcal{H}_h^p}^p = \frac{|u(x_0)|^p}{h(x_0)^p} + \int_U G_U(x_0, y) \int_{\mathbb{R}^d} F \left( \frac{u(y)}{h(y)}, \frac{u(z)}{h(z)} \right) \mathcal{A} \frac{h(y) dz dy}{h(x_0) |z-y|^{d+\alpha}}.$$

$\square$

We remark in passing that for  $h \equiv 1$  we obtain  $\mathcal{H}_h^p = \mathcal{H}^p$ . To rigorously state this observation, one should discuss conditioning by functions  $h$  with

nontrivial values on  $D^c$  ([12]). We will not embark on this endeavor, instead in the next section we will fully discuss the conditional Hardy spaces of a local operator, in which case the values of  $h$  on  $D^c$  are irrelevant.

#### 4. CLASSICAL HARDY SPACES

We will describe the Hardy spaces and the conditional Hardy spaces of harmonic functions of the Laplacian  $\Delta = \sum_{j=1}^d \partial^2/\partial x_j^2$ . The former case has been widely studied in the literature, mainly for the ball and the half-space, but also for smooth and Lipschitz domains, see [1, 25, 36, 37, 23]. The characterization of the Hardy spaces in terms of quadratic functions appeared in [35] and [43] for harmonic functions on the half-space in  $\mathbb{R}^d$ . The case of  $D$  being the unit ball was studied in detail in [40, 32]. For more general domains in  $\mathbb{R}^d$  see [41, 37, 23].

Throughout this section we will additionally assume that  $D$  is connected, i.e. it is a domain. For  $0 < p < \infty$ , the classical Hardy space  $H^p(D)$  may be defined as the family of all those functions  $u$  on  $D$  which are harmonic on  $D$  (i.e.  $u \in C^2(D)$  and  $\Delta u(x) = 0$  for  $x \in D$ ) and satisfy

$$\|u\|_{H^p} := \sup_{U \subset \subset D} \left( \mathbb{E}_{x_0} |u(W_{\tau_U})|^p \right)^{1/p} < \infty.$$

Here  $W$  is the Brownian motion on  $\mathbb{R}^d$  and  $\tau_U = \inf\{t \geq 0 : W_t \notin U\}$ . For a positive harmonic function  $h$  on  $D$  and  $0 < p < \infty$  we will consider the space  $H_h^p(D)$  of all those functions  $u$  harmonic on  $D$  which satisfy

$$\|u\|_{H_h^p}^p := \sup_{U \subset \subset D} \mathbb{E}_{x_0}^h \left| \frac{u(W_{\tau_U})}{h(W_{\tau_U})} \right|^p = \frac{1}{h(x_0)} \sup_{U \subset \subset D} \mathbb{E}_{x_0} \frac{|u(W_{\tau_U})|^p}{h(W_{\tau_U})^{p-1}} < \infty,$$

where  $\mathbb{E}_x^h$  is the expectation for the conditional Brownian motion (compare Section 3 or see [16]). Let  $G_D$  be the classical Green function of  $D$  for  $\Delta$ . If  $1 < p < \infty$  and  $u$  is harmonic on  $D$ , then the following Hardy-Stein identity holds

$$(36) \quad \|u\|_{H^p}^p = |u(x_0)|^p + p(p-1) \int_D G_D(x_0, y) |u(y)|^{p-2} |\nabla u(y)|^2 dy.$$

The identity (36) obtains by taking  $h \equiv 1$  in the next theorem. (36) generalizes [40, Lemma 1] and [32, Theorem 4.3], where the formula was given for the ball in  $\mathbb{R}^d$ , see also [38]. We note that (36) is implicit in [41, Lemma 6], but apparently the identity did not receive enough attention for general domains. We also note that in many cases the exact asymptotics of  $G_D$  is known, and we obtain explicit estimates. For instance, if  $D$  is a bounded  $C^{1,1}$  domain in  $\mathbb{R}^d$  and  $d \geq 3$ , then  $G_D(x_0, y) \asymp \delta_D(y) |y - x_0|^{2-d}$ , where  $\delta_D(y) := \text{dist}(y, D^c)$ , see [6], also for the case of Lipschitz domains.

**Theorem 3.** *If  $1 < p < \infty$  and  $u$  is harmonic on  $D$ , then*

$$(37) \quad \|u\|_{H_h^p}^p = \frac{|u(x_0)|^p}{h(x_0)^p} + p(p-1) \int_D G_D(x_0, y) \left| \frac{u(y)}{h(y)} \right|^{p-2} \left| \nabla \frac{u}{h}(y) \right|^2 \frac{h(y)}{h(x_0)} dy.$$

The remainder of this section is devoted to the proof of Theorem 3. The reader interested mostly in (36) is encouraged to carry out similar but simpler calculations for  $h \equiv 1$  and  $p > 2$ . We note that (37) is quite more general than (36) because usually  $u/h$  is not harmonic. We start with the following well-known Green-type equality. Consider open  $U \subset\subset D$  and a real-valued function  $\phi : \mathbb{R}^d \rightarrow \mathbb{R}$  which is  $C^2$  in a neighborhood of  $\bar{U}$ . Then  $\Delta\phi$  is bounded on  $\bar{U}$ , and for every  $x \in D$  we have

$$(38) \quad \phi(x) = \mathbb{E}_x \phi(W_{\tau_U}) - \int_U G_U(x, y) \Delta\phi(y) dy,$$

see, e.g., [18, p. 133] for the proof.

**Lemma 9.** *Let  $\varepsilon \neq 0$  and let  $u$  be harmonic on  $D$ . We have*

$$(39) \quad \Delta \left[ \left( \frac{u^2}{h^2} + \varepsilon^2 \right)^{p/2} h \right] = p \left( \frac{u^2}{h^2} + \varepsilon^2 \right)^{(p-4)/2} \left[ (p-1) \frac{u^2}{h^2} + \varepsilon^2 \right] \left| \nabla \frac{u}{h} \right|^2.$$

*If  $u \neq 0$  or  $p \geq 2$ , then*

$$(40) \quad \Delta \left( \frac{|u|^p}{h^{p-1}} \right) = p(p-1) \left| \frac{u}{h} \right|^{p-2} \left| \nabla \frac{u}{h} \right|^2 h.$$

*Proof.* Denote  $u_i = \partial u / \partial x_i$ ,  $h_i = \partial h / \partial x_i$ ,  $u_{ii} = \partial^2 u / \partial x_i^2$  and  $h_{ii} = \partial^2 h / \partial x_i^2$ ,  $i = 1, \dots, d$ . The lemma results from straightforward calculations based on the following observations:

$$\begin{aligned} \nabla |u|^p &= \nabla (u^2)^{p/2} = p|u|^{p-2} u \nabla u, \quad \text{if } p \geq 2 \text{ or } u \neq 0, \\ \frac{\partial^2}{\partial x_i^2} |u|^p &= p(p-1)|u|^{p-2} u_i^2 + p|u|^{p-2} u u_{ii}, \quad \text{if } p \geq 2 \text{ or } u \neq 0, \\ \nabla h^{1-p} &= (1-p) h^{-p} \nabla h, \\ \Delta(fg) &= f \Delta g + 2 \nabla f \circ \nabla g + g \Delta f. \end{aligned}$$

This yields (40) if  $p \geq 2$  or  $u(x) \neq 0$  at the point  $x$  where the derivatives are calculated (and so  $|u|^p h^{1-p}$  is of class  $C^2$  there). To prove (39) we let  $\varepsilon \neq 0$ ,

denote  $f(x) = u^2/h^2 + \varepsilon^2$ , and use a few more identities:

$$\begin{aligned}\nabla \frac{u}{h} &= \frac{\nabla u}{h} - \frac{u \nabla h}{h^2}, & \nabla \left( \frac{u}{h} \right)^2 &= 2 \frac{u}{h} \nabla \frac{u}{h}, \\ \Delta \left( \frac{u}{h} \right)^2 &= \frac{2|\nabla u|^2}{h^2} - \frac{8u \nabla u \circ \nabla h}{h^3} + \frac{6u^2 |\nabla h|^2}{h^4}, \\ \nabla f^{p/2} &= \frac{p}{2} f^{p/2-1} \nabla \left( \frac{u}{h} \right)^2, \\ \Delta f^{p/2} &= \frac{p(p-2)}{4} f^{p/2-2} \left| \nabla \left( \frac{u}{h} \right)^2 \right|^2 + \frac{p}{2} f^{p/2-1} \Delta \left( \frac{u}{h} \right)^2, \\ \Delta \left( f^{p/2} h \right) &= \frac{p(p-2)}{4} f^{p/2-2} \left| \nabla \left( \frac{u}{h} \right)^2 \right|^2 h + p f^{p/2-1} \left| \nabla \frac{u}{h} \right|^2 h.\end{aligned}$$

□

Noteworthy, we obtained nonnegative expressions in (39) and (40). Also, if  $\varepsilon \rightarrow 0$ , then  $\Delta[(u^2/h^2 + \varepsilon^2)^{p/2} h] \rightarrow \Delta(|u|^p h^{1-p})$  almost everywhere on  $D$ .

**Lemma 10.** *If  $u$  is harmonic on  $D$ ,  $U \subset\subset D$  and  $p > 1$ , then*

$$\mathbb{E}_{x_0} \frac{|u(X_{\tau_U})|^p}{h(X_{\tau_U})^{p-1}} = \frac{|u(x_0)|^p}{h(x_0)^{p-1}} + p(p-1) \int_U G_U(x_0, y) \left| \frac{u(y)}{h(y)} \right|^{p-2} \left| \nabla \frac{u}{h}(y) \right|^2 h(y) dy.$$

*Proof.* For  $p \geq 2$  we have  $|u|^p h^{1-p} \in C^2(D)$  and the result follows from (38) and Lemma 9. If  $1 < p < 2$ , then we consider  $u + i\varepsilon h$  in place of  $u$  and we let  $\varepsilon \rightarrow 0$ . By (39), (38), Fatou's lemma and dominated convergence we obtain the result. □

*Proof of Theorem 3.* The result follows from Lemma 10 and monotone convergence, after dividing by  $h(x_0)$ . □

In view of close similarities between the above descriptions of Hardy and conditional Hardy spaces, it seems that the framework of conditional semi-groups (23) is appropriate for such calculations in more general settings.

## 5. FURTHER RESULTS

We will now discuss the structure of  $\mathcal{H}^p$ . We start with  $p = 1$ . The following is a counterpart of the theorem of Krickeberg for martingales ([15]), and an extension of [30, Theorem 1], where the result was proved for singular  $\alpha$ -harmonic functions on bounded Lipschitz domains.

**Lemma 11.** *Let  $u \in \mathcal{H}^1$ . There exist nonnegative functions  $f$  and  $g$  which are  $\alpha$ -harmonic on  $D$ , satisfy  $u = f - g$  and uniquely minimize  $f(x_0) + g(x_0)$ .*

In fact,  $f(x_0) + g(x_0) = \|u\|_1$ . If  $u$  is singular  $\alpha$ -harmonic on  $D$ , then so are  $f$  and  $g$ . If  $1 \leq p < \infty$  and  $u \in \mathcal{H}^p$ , then  $\|u\|_p^p = \|f\|_p^p + \|g\|_p^p$ .

*Proof.* Let  $U_n$  be open,  $U_n \subset\subset U_{n+1}$  for  $n = 1, 2, \dots$  and  $\bigcup_n U_n = D$ . Let  $\tau_n = \tau_{U_n}$ . We have

$$\|u\|_1 = \lim_{n \rightarrow \infty} \mathbb{E}_{x_0} |u(X_{\tau_n})| < \infty.$$

Let  $u^+ = \max(u, 0)$  and  $u^- = \max(-u, 0)$ . For  $n = 1, 2, \dots$ , we define

$$f_n(x) = \mathbb{E}_x u^+(X_{\tau_n}), \quad g_n(x) = \mathbb{E}_x u^-(X_{\tau_n}), \quad x \in \mathbb{R}^d.$$

Obviously, functions  $f_n$  and  $g_n$  are nonnegative on  $\mathbb{R}^d$ , and finite and  $\alpha$ -harmonic on  $U_n$ . We have  $u = f_n - g_n$ . Since  $\tau_n \leq \tau_{n+1}$ , for every  $x \in \mathbb{R}^d$ ,

$$f_n(x) = \mathbb{E}_x [\mathbb{E}_{X_{\tau_n}} u(X_{\tau_{n+1}}) ; u(X_{\tau_n}) > 0] \leq \mathbb{E}_x [\mathbb{E}_{X_{\tau_n}} u^+(X_{\tau_{n+1}})] = f_{n+1}(x),$$

and  $g_n(x) \leq g_{n+1}$ . We let  $f(x) = \lim f_n(x)$  and  $g(x) = \lim g_n(x)$ . By the monotone convergence theorem, the mean value property (2) holds for  $f$  and  $g$ . We obtain

$$f(x_0) + g(x_0) = \lim_{n \rightarrow \infty} \mathbb{E}_{x_0} |u(X_{\tau_n})| = \|u\|_1 < \infty.$$

In view of Harnack inequality we conclude that  $f$  and  $g$  are finite, hence  $\alpha$ -harmonic on  $D$ . Also,  $u = f - g$ . If  $u$  vanishes on  $D$ , then so do  $f$  and  $g$ . For the uniqueness, we observe that if  $\tilde{f}, \tilde{g} \geq 0$  are  $\alpha$ -harmonic on  $D$ , and  $u = \tilde{f} - \tilde{g}$ , then  $-\tilde{g} \leq u \leq \tilde{f}$ , hence  $\tilde{f} \leq f$  and  $g \leq \tilde{g}$  by the construction of  $f$  and  $g$ . Therefore  $f(x_0) + g(x_0) \leq \tilde{f}(x_0) + \tilde{g}(x_0)$ , and equality holds if and only if  $f(x_0) = \tilde{f}(x_0)$  and  $g(x_0) = \tilde{g}(x_0)$ , henceforth  $f = \tilde{f}$  and  $g = \tilde{g}$ .

Let  $p > 1$  and suppose that  $u \in \mathcal{H}^p \subset \mathcal{H}^1$ . By Jensen's inequality,

$$f_n(x)^p \leq \mathbb{E}_x u^+(X_{\tau_n})^p, \quad g_n(x)^p \leq \mathbb{E}_x u^-(X_{\tau_n})^p,$$

hence

$$f_n(x)^p + g_n(x)^p \leq \mathbb{E}_x |u(X_{\tau_n})|^p.$$

For  $m < n$  we have

$$\mathbb{E}_{x_0} (f_n(X_{\tau_m})^p + g_n(X_{\tau_m})^p) \leq \mathbb{E}_{x_0} \mathbb{E}_{X_{\tau_m}} |u(X_{\tau_n})|^p = \mathbb{E}_{x_0} |u(X_{\tau_n})|^p.$$

Letting  $n \rightarrow \infty$ , we get

$$\mathbb{E}_{x_0} (f(X_{\tau_m})^p + g(X_{\tau_m})^p) \leq \|u\|_p^p.$$

Hence  $\|f\|_p^p + \|g\|_p^p \leq \|u\|_p^p$ . On the other hand,  $f, g \geq 0$ , hence

$$\begin{aligned} \|u\|_p^p &= \lim_{n \rightarrow \infty} \mathbb{E}_{x_0} |f(X_{\tau_n}) - g(X_{\tau_n})|^p \\ &\leq \lim_{n \rightarrow \infty} \mathbb{E}_{x_0} (f(X_{\tau_n})^p + g(X_{\tau_n})^p) = \|f\|_p^p + \|g\|_p^p. \end{aligned}$$

The proof is complete.  $\square$

We note that  $\|u\|_p^p = \|f\|_p^p + \|g\|_p^p$  has a trivial analogue for  $L^p$  spaces.

In view of Lemma 11 and the representation of positive  $\alpha$ -harmonic functions [12, Theorem 3], each  $u \in \mathcal{H}^1$  has a unique integral representation

$$(41) \quad u(x) = \int_{\partial_M D} M_D(x, z) \mu(dz) + \int_{D^c} f(y) P_D(x, y) dy, \quad x \in D,$$

where  $\mu$  is a (finite) signed measure on the Martin boundary of  $D$ ,  $M_D$  is the Martin kernel of  $D$ ,  $P_D$  is the Poisson kernel of  $D$ , and the function  $f$  satisfies  $\int_{D^c} |f(y)| P_D(x_0, y) dy < \infty$ , see [12].

**Lemma 12.** *Let  $u \in \mathcal{H}_h^1$ . There are nonnegative functions  $f, g \in \mathcal{H}_h^1$  which satisfy  $u = f - g$  and uniquely minimize  $f(x_0) + g(x_0)$ . In fact,  $f(x_0) + g(x_0) = \|u\|_{\mathcal{H}_h^1} h(x_0)$ . If  $1 \leq p < \infty$  and  $u \in \mathcal{H}_h^p$ , then  $\|u\|_{\mathcal{H}_h^p}^p = \|f\|_{\mathcal{H}_h^p}^p + \|g\|_{\mathcal{H}_h^p}^p$ .*

*Proof.* If  $u \in \mathcal{H}_h^1$ , then  $u$  is singular  $\alpha$ -harmonic on  $D$ ,  $u \in \mathcal{H}^1$  and  $\|u\|_{\mathcal{H}_h^1} = h(x_0)^{-1} \|u\|_1$  (conditioning is trivial for  $p = 1$ ). By Lemma 11,  $u$  has the Krickeberg decomposition  $u = f - g$ , and  $f, g$  are nonnegative and singular  $\alpha$ -harmonic on  $D$ . In particular  $\|f\|_{\mathcal{H}_h^1} = f(x_0)/h(x_0)$  and  $\|g\|_{\mathcal{H}_h^1} = g(x_0)/h(x_0)$  are finite. The reader may easily verify the rest of the statement of the lemma, following the previous proof and using the conditional expectation  $\mathbb{E}^h$ .  $\square$

**Remark 2.** *Analogues of Lemma 11 and Lemma 12 are true for the classical Hardy spaces  $H^p(D)$  and  $H_h^p(D)$  for connected  $D$ .*

As an application of (36) we give a short proof of the following Littlewood-Paley type inequality (see [31], where the result was given for the ball in  $\mathbb{R}^2$ ). Recall the notation  $\delta_D(y) = \text{dist}(y, D^c)$ .

**Proposition 13.** *Consider a domain  $D \subset \mathbb{R}^d$ , and let  $p \geq 2$ . There is a constant  $c > 0$  such that for every function  $u$  harmonic on  $D$ ,*

$$\|u\|_{H^p}^p - |u(x_0)|^p \geq p(p-1)d^{2-p}2^{1-p} \int_D G_D(x_0, y) \delta_D(y)^{p-2} |\nabla u(y)|^p dy.$$

*Proof.* We may assume that  $\|u\|_{H^p} < \infty$ . In view of Lemma 11 and Remark 2,  $u = f - g$ , where  $f, g$  are positive and harmonic on  $D$  and  $\|u\|_{H^p}^p = \|f\|_{H^p}^p + \|g\|_{H^p}^p$ . Clearly,  $|u(x_0)|^p \leq f(x_0)^p + g(x_0)^p$ , hence  $\|u\|_{H^p}^p - |u(x_0)|^p \geq \|f\|_{H^p}^p - |f(x_0)|^p + \|g\|_{H^p}^p - |g(x_0)|^p$ . Furthermore, by Jensen's inequality,

$$|\nabla u|^p \leq 2^{p-1} (|\nabla f|^p + |\nabla g|^p).$$

Recall the following gradient estimate for the nonnegative harmonic function  $f$ ,

$$f(x) \geq |\nabla f(x)| \delta_D(x) / d, \quad x \in D,$$

([20, Exercise 2.13], see also [2]). Here  $d$  is the dimension. By (36),

$$\begin{aligned} \|f\|_{H^p}^p - |f(x_0)|^p &= p(p-1) \int_D G_D(x_0, y) |f(y)|^{p-2} |\nabla f(y)|^2 dy \\ &\geq p(p-1) d^{2-p} \int_D G_D(x_0, y) \delta_D(y)^{p-2} |\nabla f(y)|^p dy, \end{aligned}$$

and a similar estimate holds for  $g$ .  $\square$

**Acknowledgements.** Part of the above results on  $\mathcal{H}^p(D, \alpha)$  were presented at the 4th International Conference on Stochastic Analysis and Its Applications held at Kansai University, Osaka, August 30 - September 3, 2010. We gratefully thank the organizers for the opportunity.

#### REFERENCES

- [1] Sheldon Axler, Paul Bourdon, and Wade Ramey. *Harmonic function theory*, volume 137 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 2001.
- [2] Rodrigo Bañuelos and Michael M.H. Pang. Lower bound gradient estimates for solutions of Schrödinger equations and heat kernels. *Comm. Partial Differential Equations*, 24(3-4):499–543, 1999.
- [3] Richard F. Bass and Dahae You. A Fatou theorem for  $\alpha$ -harmonic functions in Lipschitz domains. *Probab. Theory Related Fields*, 133(3):391–408, 2005.
- [4] Jürgen Bliedtner and Wolfhard Hansen. *Potential theory*. Universitext. Springer-Verlag, Berlin, 1986. An analytic and probabilistic approach to balayage.
- [5] Krzysztof Bogdan. The boundary Harnack principle for the fractional Laplacian. *Studia Math.*, 123(1):43–80, 1997.
- [6] Krzysztof Bogdan. Sharp estimates for the Green function in Lipschitz domains. *J. Math. Anal. Appl.*, 243(2):326–337, 2000.
- [7] Krzysztof Bogdan, Krzysztof Burdzy, and Zhen-Qing Chen. Censored stable processes. *Probab. Theory Related Fields*, 127(1):89–152, 2003.
- [8] Krzysztof Bogdan and Tomasz Byczkowski. Potential theory for the  $\alpha$ -stable Schrödinger operator on bounded Lipschitz domains. *Studia Math.*, 133(1):53–92, 1999.
- [9] Krzysztof Bogdan, Tomasz Byczkowski, Tadeusz Kulczycki, Michal Ryznar, Renming Song, and Zoran Vondraček. *Potential analysis of stable processes and its extensions*, volume 1980 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2009. Edited by Piotr Graczyk and Andrzej Stos.
- [10] Krzysztof Bogdan and Bartłomiej Dyda. Relative Fatou theorem for harmonic functions of rotation invariant stable processes in smooth domains. *Studia Math.*, 157(1):83–96, 2003.
- [11] Krzysztof Bogdan and Tomasz Jakubowski. Estimates of the Green function for the fractional Laplacian perturbed by gradient. *Potential Analysis*, pages 1–27. 10.1007/s11118-011-9237-x.
- [12] Krzysztof Bogdan, Tadeusz Kulczycki, and Mateusz Kwaśnicki. Estimates and structure of  $\alpha$ -harmonic functions. *Probab. Theory Related Fields*, 140(3-4):345–381, 2008.
- [13] Krzysztof Bogdan and Paweł Sztonyk. Estimates of the potential kernel and Harnack’s inequality for the anisotropic fractional Laplacian. *Studia Math.*, 181(2):101–123, 2007.

- [14] Krzysztof Bogdan and Tomasz Żak. On Kelvin transformation. *J. Theoret. Probab.*, 19(1):89–120, 2006.
- [15] Claude Dellacherie and Paul-André Meyer. *Probabilities and potential. B*, volume 72 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam, 1982. Theory of martingales, Translated from the French by J. P. Wilson.
- [16] Joseph L. Doob. *Classical Potential Theory and Its Probabilistic Counterpart*, volume 262 of *Die Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, New York, 1984.
- [17] Richard Durrett. *Brownian motion and martingales in analysis*. Wadsworth Mathematics Series. Wadsworth International Group, Belmont, CA, 1984.
- [18] Eugene B. Dynkin. *Markov Processes, I*, volume 121 of *Die Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin-Göttingen-Heidelberg, 1965. Translated with the authorization and assistance of the author by J. Fabius, V. Greenberg, A. Maitra, G. Majone.
- [19] Masatoshi Fukushima, Yōichi Ōshima, and Masayoshi Takeda. *Dirichlet forms and symmetric Markov processes*, volume 19 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, 1994.
- [20] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*, volume 224 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 1983.
- [21] Qing-Yang Guan and Zhi-Ming Ma. Reflected symmetric  $\alpha$ -stable processes and regional fractional Laplacian. *Probab. Theory Related Fields*, 134(4):649–694, 2006.
- [22] Tomasz Jakubowski. The estimates for the Green function in Lipschitz domains for the symmetric stable processes. *Probab. Math. Statist.*, 22(2, Acta Univ. Wratislav. No. 2470):419–441, 2002.
- [23] David S. Jerison and Carlos E. Kenig. Boundary value problems on Lipschitz domains. In *Studies in partial differential equations*, volume 23 of *MAA Stud. Math.*, pages 1–68. Math. Assoc. America, Washington, DC, 1982.
- [24] Panki Kim. Relative Fatou’s theorem for  $(-\Delta)^{\alpha/2}$ -harmonic functions in bounded  $\kappa$ -fat open sets. *J. Funct. Anal.*, 234(1):70–105, 2006.
- [25] Paul Koosis. *Introduction to  $H_p$  Spaces*, volume 115 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998. Second edition.
- [26] Naum S. Landkof. *Foundations of modern potential theory*. Springer-Verlag, New York, 1972. Translated from the Russian by A. P. Doohovskoy, Die Grundlehren der mathematischen Wissenschaften, Band 180.
- [27] Tomasz Luks. Hardy spaces for the Laplacian with lower order perturbations. *Studia Math.*, 204:39–62, 2011.
- [28] Olivier Mazet. A characterization of Markov property for semigroups with invariant measure. *Potential Anal.*, 16(3):279–287, 2002.
- [29] Krzysztof Michalik and Michał Ryznar. Relative Fatou theorem for  $\alpha$ -harmonic functions in Lipschitz domains. *Illinois J. Math.*, 48(3):977–998, 2004.
- [30] Krzysztof Michalik and Michał Ryznar. Hardy spaces for  $\alpha$ -harmonic functions in regular domains. *Math. Z.*, 265(1):173–186, 2010.
- [31] Miroslav Pavlović. A short proof of an inequality of Littlewood and Paley. *Proc. Amer. Math. Soc.*, 134(12):3625–3627, 2006.
- [32] Miroslav Pavlović. Green’s formula and the Hardy-Stein identities. *Filomat*, 23(3):135–153, 2009.

- [33] Miroslav Pavlović. Hardy-Stein type characterization of harmonic Bergman spaces. *Potential Anal.*, 32(1):1–15, 2010.
- [34] Philip E. Protter. *Stochastic integration and differential equations*, volume 21 of *Stochastic Modelling and Applied Probability*. Springer-Verlag, Berlin, 2005. Second edition. Version 2.1, Corrected third printing.
- [35] Elias M. Stein. On the theory of harmonic functions of several variables. II. Behavior near the boundary. *Acta Math.*, 106:137–174, 1961.
- [36] Elias M. Stein. *Singular integrals and differentiability properties of functions*. Princeton Mathematical Series, No. 30. Princeton University Press, Princeton, N.J., 1970.
- [37] Elias M. Stein. *Boundary behavior of holomorphic functions of several complex variables*. Princeton University Press, Princeton, N.J., 1972. Mathematical Notes, No. 11.
- [38] Peter Stein. On a theorem of M. Riesz. *J. Lond. Math. Soc.*, 8:242–247, 1933.
- [39] Stevo Stević. On harmonic Hardy and Bergman spaces. *J. Math. Soc. Japan*, 54(4):983–996, 2002.
- [40] Stevo Stević. On harmonic Hardy spaces and area integrals. *J. Math. Soc. Japan*, 56(2):339–347, 2004.
- [41] Manfred Stoll. The Littlewood-Paley inequalities for Hardy-Orlicz spaces of harmonic functions on domains in  $\mathbb{R}^n$ . In *Potential theory in Matsue*, volume 44 of *Adv. Stud. Pure Math.*, pages 363–376. Math. Soc. Japan, Tokyo, 2006.
- [42] Ferenc Weisz. *Martingale Hardy spaces and their applications in Fourier analysis*, volume 1568 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1994.
- [43] Kjell-Ove Widman. On the boundary behavior of solutions to a class of elliptic partial differential equations. *Ark. Mat.*, 6(6):485–533, 1967.

INSTITUTE OF MATHEMATICS AND COMPUTER SCIENCE, WROCLAW UNIVERSITY OF TECHNOLOGY, WYBRZEŻE WYSPIAŃSKIEGO 27, 50-370 WROCLAW, POLAND  
*E-mail address:* Krzysztof.Bogdan@pwr.wroc.pl

FACULTY OF MATHEMATICS, UNIVERSITY OF BIELEFELD, POSTFACH 10 01 31, D-33501 BIELEFELD, GERMANY AND INSTITUTE OF MATHEMATICS AND COMPUTER SCIENCE, WROCLAW UNIVERSITY OF TECHNOLOGY, WYBRZEŻE WYSPIAŃSKIEGO 27, 50-370 WROCLAW, POLAND  
*E-mail address:* bdyda@pwr.wroc.pl

LAREMA, LABORATOIRE DE MATHÉMATIQUES, UNIVERSITÉ D'ANGERS, 2 BOULEVARD LAVOISIER, 49045 ANGERS CEDEX 01, FRANCE  
*E-mail address:* luks@math.univ-angers.fr