

A FRAMEWORK FOR FRACTIONAL HARDY INEQUALITIES

BARTŁOMIEJ DYDA AND ANTTI V. VÄHÄKANGAS

ABSTRACT. We provide a general framework for fractional Hardy inequalities. Our framework covers, for instance, fractional inequalities related to the Dirichlet forms of some Lévy processes, and weighted fractional inequalities on irregular open sets.

1. INTRODUCTION

The objective of the present paper is to study inequalities of the general form

$$(1.1) \quad \int_D \frac{|u(x)|^p}{\phi(\delta_x)} \mu(dx) \leq c \int_D \int_{D \cap B(x, R\delta_x)} \frac{|u(x) - u(y)|^p}{\phi(\delta_x) \delta_x^d} \mu(dy) \mu(dx), \quad c, R > 0,$$

on metric measure spaces (X, ρ, μ) and partly on $X = \mathbb{R}^d$ equipped with the Euclidean distance and the Lebesgue measure. We write $\delta_x = \text{dist}(x, X \setminus D)$ and $D \subset X$ is a possibly irregular open set. The function $\phi : (0, \infty) \rightarrow (0, \infty)$ is a ‘perturbation’ of a power function $t \mapsto t^\eta$ for some $\eta \in \mathbb{R}$, and the exponent p satisfies $0 < p < \infty$. Our main result, Theorem 4 in §3, brings together two so-far distinct lines along which the fractional Hardy inequality has been generalised: one of them related to the function ϕ , and the other to the regularity of the open set $D \subset X$. Although Theorem 4, allows for both general open sets D and functions ϕ simultaneously, we will nevertheless state separate results in each of these directions to make the exposition simpler.

Regularly varying functions ϕ with index $\eta > 0$ are allowed in (1.1), see §3.1 and [3]. Let us remind that ϕ is called *regularly varying* at origin (resp. infinity) with index η , if $\frac{\phi(\lambda x)}{\phi(x)} \rightarrow \lambda^\eta$ when $x \rightarrow 0_+$ (resp. when $x \rightarrow \infty$) for every $\lambda > 0$. We state the following theorem as an example.

Theorem 1. *Suppose that $\phi : (0, \infty) \rightarrow (0, \infty)$ is a regularly varying function at origin of index ρ_0 , a regularly varying function at infinity of index ρ_∞ , and is bounded and bounded away from zero on every compact subset of $(0, \infty)$. Suppose that either $0 < \rho_0, \rho_\infty < d$ or $\rho_0, \rho_\infty > d$. Let $p > 0$. Then there exists a constant $c = c(\phi, d, p)$ such that*

$$(1.2) \quad \int_{\mathbb{R}^d \setminus \{0\}} \frac{|u(x)|^p}{\phi(|x|)} dx \leq c \int_{\mathbb{R}^d \setminus \{0\}} \int_{\mathbb{R}^d \setminus \{0\}} \frac{|u(x) - u(y)|^p}{\phi(|x - y|) |x - y|^d} dy dx$$

for every measurable function u for which the left hand side is finite.

2010 *Mathematics Subject Classification.* 26D15, 46E35, 31C25, 26A12.

Key words and phrases. fractional Hardy inequality, fractional Laplacian, regularly varying function, Assouad dimension, uniform fatness, irregular domain.

Inequalities like (1.2) have been studied in [15] for weights of more general (but also more complicated) form and $p > 1$, and in [14, 15, 22] in the one-dimensional case. The forms appearing on the right hand side of (1.2) for $p = 2$ (and for more general domains) are, at least for some functions ϕ , the Dirichlet forms of some Lévy processes, which are being extensively studied, see e.g. [13, 20, 33] and [5, Section 4.1] for examples.

To discuss our results for irregular open sets, we confine ourselves to weighted fractional Hardy inequalities in \mathbb{R}^d , i.e., we consider a function $\phi(t) = t^{sp-\beta}$ with $d + sp \geq 0$, in which case inequality (1.1) yields

$$(1.3) \quad \int_D \frac{|u(x)|^p}{\delta_x^{sp}} \delta_x^\beta dx \leq cR^{d+sp} \int_D \int_D \frac{|u(x) - u(y)|^p}{|x - y|^{d+sp}} \delta_x^\beta dy dx.$$

An open set $D \subset \mathbb{R}^d$ is said to admit (s, p, β) -Hardy inequality, if inequality (1.3) holds for all functions $u \in C_0^\infty(D)$ (i.e., smooth with compact support in D) with $c, R > 0$ independent of u . An analogous weighted (p, β) -Hardy inequality, with $c > 0$,

$$(1.4) \quad \int_D \frac{|u(x)|^p}{\delta_x^p} \delta_x^\beta dx \leq c \int_D |\nabla u(x)|^p \delta_x^\beta dx$$

holds for every $u \in C_0^\infty(D)$ if D is a bounded Lipschitz domain, $1 < p < \infty$, and $\beta < p - 1$, [32]. More generally, an open set admits a (p, β) -Hardy inequality if the complement $D^c = \mathbb{R}^d \setminus D$ is either sufficiently ‘thin’ or ‘fat’. For instance, an open set D admits a $(p, 0)$ -Hardy inequality if D^c is $(1, p)$ -uniformly fat and $1 < p < \infty$, [26]. The $(1, p)$ -fatness of D^c is also known to be sufficient for certain (p, β) -Hardy inequalities, we refer to [24, 37]. A deeper understanding of the dichotomy is reached in an independent recent study [23], where an open set $D \subset X$ is shown to admit a (p, β) -Hardy inequality if $D^c = X \setminus D$ sufficiently thin or fat, measured in terms of upper and lower Assouad dimension ($\overline{\dim}_A$ and $\underline{\dim}_A$), respectively. We also refer to [21].

Our framework covers an Assouad dichotomy result for fractional (s, p, β) -Hardy inequalities with $X = \mathbb{R}^d$, see Theorem 2. As a matter of fact, it covers more general fractional inequalities of Hardy-type under similar geometric assumptions. The simple proof of our main result is a refinement of techniques in [7] where, e.g., $(s, p, 0)$ -Hardy inequalities for bounded Lipschitz domains are found. There has been recent interest in $(s, p, 0)$ -Hardy inequalities and the boundary regularity of an open set D , we refer to [10, 16, 17, 18]. In another direction, the sharp constants for fractional Hardy-type inequalities on general domains are obtained in [27], where the distance is replaced with an averaged pseudo distance. In [9] these results are refined and other related papers are [4, 8, 11, 12, 34].

The following is our Assouad dichotomy result. For the relevant definitions, we refer to §2.

Theorem 2. *Let p, s, β be real numbers so that $0 < p < \infty$ and $d + sp \geq 0$. Suppose $D \neq \emptyset$ is a proper κ -plump open set in \mathbb{R}^d so that either condition (T) or condition (F) holds.*

(T) $\overline{\dim}_A(\partial D) < d - sp + \beta$ and D is unbounded;

(F) $\underline{\dim}_A(\partial D) > d - sp + \beta$ and, if D is unbounded, then ∂D is unbounded.

Then D admits an (s, p, β) -Hardy inequality.

This theorem follows from Theorem 4 with the aid of propositions 5 and 9. As an illustrative example, we may consider the Koch snowflake domain $D \subset \mathbb{R}^2$. It is a bounded domain with a property $\underline{\dim}_A(\partial D) = \log 4 / \log 3$. Hence, Theorem 2 does apply since D is also κ -plump. In a ‘thin case’ we may, e.g., consider the domain $G := \mathbb{R}^d \setminus \overline{D}$, where D is the Koch snowflake. Now G is κ -plump and it satisfies $\overline{\dim}_A(\partial G) = \log 4 / \log 3$.

We comment on the cases (T) and (F) in Theorem 2. Focusing on the case (T) first, recall that $\overline{\dim}_A(\partial D) = d - 1$ for a Lipschitz domain D . The unboundedness of D cannot be removed, at least if $0 < s < 1$, in which case a bounded Lipschitz domain satisfies an $(s, p, 0)$ -Hardy inequality if and only if $sp > 1$, [7]. Certain non-homogeneous $(s, p, 0)$ -Hardy inequalities remain valid for John domains D with $\overline{\dim}_A(\partial D) < d - sp$, [18]. Therein (T) with $\beta = 0$ is formulated in terms of a certain Aikawa dimension which equals to the upper Assouad dimension in Euclidean spaces, see [25]. Recalling that John domains are both bounded and κ -plump, we may conclude that our framework provides a far-reaching generalisation of the non-homogeneous results to the case of unbounded open sets.

The dimensional restriction in (T) is somewhat natural: under some *a priori* conditions on D , the inequality $\overline{\dim}_A(\partial D) < d - sp$ is equivalent with non-homogeneous $(s, p, 0)$ -Hardy inequality on Triebel–Lizorkin spaces, [18]. Likewise, the κ -plumpness condition is natural in some cases. For illustration, let $D = \mathbb{R}^d \setminus K$, where K is a closed set such that $\partial D = K$. Now the κ -plumpness of $D = \mathbb{R}^d \setminus K$ with some $\kappa \in (0, 1)$ is characterised by inequality (T) with $d - sp + \beta = d$, i.e.,

$$\overline{\dim}_A(K) = \overline{\dim}_A(\partial D) < d.$$

We refer to [28] for further results on so-called porosity and the upper Assouad dimension.

Moving on to the case (F) with ‘fat’ boundary, let us first formulate an illustrative, but more restrictive, corollary of Theorem 2. We refer to §4.4 for the relevant definitions.

Corollary 3. *Let p, s, β be real numbers so that $1 < p < \infty$, $0 < sp - \beta < d$, and $d + sp \geq 0$. Suppose D is a κ -plump open set in \mathbb{R}^d such that ∂D is $(s - \beta/p, p)$ -uniformly fat (-locally uniformly fat, if D is bounded). Then D admits an (s, p, β) -Hardy inequality.*

This corollary is a consequence of Theorem 2 and propositions 7 and 8. Unlike in the case of inequality (1.4) with $\beta = 0$ and $s = 1$, the (s, p) -uniform fatness of ∂D (let alone D^c) is not a sufficient condition for an open set D to admit an $(s, p, 0)$ -Hardy inequality (at least) in the case of $0 < sp \leq 1$. This ‘non-local obstruction’ is recognised and addressed in [16]. It affects certain fractional Hardy inequalities that are treated in [10]. One assumes there that D^c is (s, p) -uniformly fat and, as a conclusion, on the right hand side of (1.3) one has integration over $\mathbb{R}^d \times \mathbb{R}^d$ instead of $D \times D$.

Suppose that D is an open set whose boundary is (s, p) -uniformly fat (locally uniformly fat, if D is bounded). It is an interesting question, what additional conditions are sufficient for D to admit an $(s, p, 0)$ -Hardy inequality. To this end, we improve a corollary in [16] where uniformity (see [29, 36]) of a domain D is shown to be a sufficient additional condition. Indeed, by Corollary 3, we may replace uniformity with κ -plumpness. Let us remark that the main result in [16], stated in terms of a ‘visibility condition on the boundary’, still covers some other cases where our results do not apply, e.g., certain domains with outward cusps.

In §2 we define both the lower and upper Assouad dimension, and the notion of κ -plumpness. We also present other basic notation. Our main result is Theorem 4, stated and proven in §3. There we also define classes WLSC and WUSC of functions ϕ and a condition DC(α, γ, d) for open sets D . The latter condition is further clarified in sections 4 and 5, where we study the cases of ‘fat’ and ‘thin’ boundaries in terms of uniform fatness, and the lower and upper Assouad dimension.

Acknowledgment. Research is supported by the DFG through SFB-701 ‘Spectral Structures and Topological Methods in Mathematics’. Part of the research was done while the second author was visiting University of Bielefeld, and he would like to thank B. Dyda and M. Kaßmann for their hospitality. The authors would like to thank K. Bogdan, T. Grzywny and J. Lehrbäck for helpful discussions and preprints of [5] and [23].

2. ASSOUD DIMENSIONS AND PLUMPNESS

We recall the lower and upper Assouad dimensions of a set $\emptyset \neq E \subset \mathbb{R}^d$, [19]. The lower Assouad dimension measures the ‘fatness’ of a set E , whereas the upper one measures how ‘thin’ a set E is. The upper Assouad dimension is often called *Assouad dimension*, a notion tracing back to [2] and even [6]. We refer to [19, 28] for further information and other results.

Definition 2.1. Consider all $\lambda \geq 0$ for which there is $C > 0$ so that, if $0 < r < R < 2 \operatorname{diam}(E)$ and $x \in E$, then at least $C(R/r)^\lambda$ balls—centred in E and of radius r —are needed to cover $B(x, R) \cap E$. The supremum of all such λ is called the *lower Assouad dimension* of E and it is denoted by $\underline{\dim}_A(E)$.

Definition 2.2. Consider all $\lambda \geq 0$ for which there is $C > 0$ so that, if $0 < r < R < 2 \operatorname{diam}(E)$ and $x \in E$, then we can cover $E \cap B(x, R)$ by at most $N \leq C(R/r)^\lambda$ balls B_1, \dots, B_N such that each B_j is centred in E and has radius r . We call the infimum of all such λ the *upper Assouad dimension* of E , and write it as $\overline{\dim}_A(E)$.

We also recall a geometric notion from [36]. See also [30].

Definition 2.3. A set $A \subset \mathbb{R}^d$ is κ -*plump* with $\kappa \in (0, 1)$ if, for each $0 < r < \operatorname{diam}(A)$ and each $x \in \bar{A}$, there is $z \in \bar{B}(x, r)$ such that $B(z, \kappa r) \subset A$.

Here is other notation; (X, ρ, μ) is a metric measure space, and we denote $\delta_x = \operatorname{dist}(x, D^c)$ with $D^c = X \setminus D$. The open ball centred at $x \in X$ and of radius $r > 0$ is denoted by $B(x, r) \subset X$. The boundary of set A is written as ∂A and $|A|$ is the Lebesgue measure of a measurable set $A \subset \mathbb{R}^d$. For a proper open set $D \subset \mathbb{R}^d$, we fix its Whitney decomposition $\mathcal{W}(D)$, and write $\mathcal{W}_m(D)$ for the family of Whitney cubes with side length 2^{-m} , $m \in \mathbb{Z}$. If $Q \in \mathcal{W}(D)$, then

$$(2.1) \quad \operatorname{diam}(Q) \leq \operatorname{dist}(Q, \partial D) \leq 4 \operatorname{diam}(Q).$$

For other properties of Whitney cubes we refer to [35, VI.1].

3. MAIN RESULT

We state and prove our main result. For definition of conditions $DC(\alpha, \gamma, d)$, $WLSC(\eta, 0, H)$ and $WUSC(\eta, 0, H)$, we refer to §3.2 and §3.1. The proof of Theorem 4 is taken up in §3.3.

Theorem 4. *Suppose that a proper open set $D \subset X$ satisfies $DC(\alpha, \gamma, d)$ with $\alpha \in (0, \infty) \setminus \{1\}$. Moreover, suppose that for some $H \in (0, 1]$, either $\alpha \in (0, 1)$, $\eta + \gamma - d > 0$ and $\phi \in WLSC(\eta, 0, H)$, or $\alpha > 1$, $\eta + \gamma - d < 0$ and $\phi \in WUSC(\eta, 0, H^{-1})$. Then for any $0 < p < \infty$ there exist constants c and $R > 0$ such that*

$$(3.1) \quad \int_D \frac{|u(x)|^p}{\phi(\delta_x)} \mu(dx) \leq c \int_D \int_{D \cap B(x, R\delta_x)} \frac{|u(x) - u(y)|^p}{\phi(\delta_x) \delta_x^d} \mu(dy) \mu(dx)$$

for all measurable functions u for which the left hand side is finite.

3.1. Assumptions on a function ϕ . We adopt the notion of a global weak lower (or upper) scaling condition (WLSC or WUSC for short) from [5, Section 3]. We formulate them in an equivalent way, which is more convenient for our purposes than the original formulation.

Definition 3.1. Let $\eta \in \mathbb{R}$ and $H \in (0, 1]$. We say that a function $\phi : (0, \infty) \rightarrow (0, \infty)$ satisfies global $WLSC(\eta, 0, H)$ (resp., $WUSC(\eta, 0, H^{-1})$) and write $\phi \in WLSC(\eta, 0, H)$ ($\phi \in WUSC(\eta, 0, H^{-1})$), if

$$(3.2) \quad \phi(st) \geq Ht^\eta \phi(s), \quad s > 0,$$

for every $t \geq 1$ (resp., for every $t \in (0, 1]$).

Remark 3.2. If the domain D in Theorem 4 is bounded, then it suffices to assume (3.2) for all $s, st < \text{diam}(D)$.

Example 3.3. Function $\phi(x) = x^\eta$, $\eta \in \mathbb{R}$, satisfies $WLSC(\eta, 0, 1)$ and $WUSC(\eta, 0, 1)$.

Example 3.4. Suppose that $\phi : (0, \infty) \rightarrow (0, \infty)$ is a regularly varying function at origin of index ρ_0 , a regularly varying function at infinity of index ρ_∞ , and is bounded and bounded away from zero on every compact subset of $(0, \infty)$. If $\rho_0 > \eta$ and $\rho_\infty > \eta$, then $\phi \in WLSC(\eta, 0, H)$ for some $H \in (0, 1]$, and if $\rho_0 < \eta$ and $\rho_\infty < \eta$, then $\phi \in WUSC(\eta, 0, H^{-1})$ for some $H \in (0, 1]$. These follow from Potter's theorem [3, Theorem 1.5.6].

We note that if, say, $\alpha < 1$, $\rho_0 \geq \eta$, $\rho_\infty \geq \eta$, and if $\eta + \gamma - d > 0$ and the assumptions on domain in Theorem 4 hold, then also the assertion (3.1) holds. Indeed, for every $\varepsilon > 0$ function ϕ satisfies $WLSC(\eta - \varepsilon, 0, H_\varepsilon)$ with some constant $H_\varepsilon \in (0, 1]$, hence by taking $\varepsilon > 0$ small enough we still have $(\eta - \varepsilon) + \gamma - d > 0$.

To have more concrete examples, let us note that functions

$$\phi_1(x) = x^\alpha + x^\beta,$$

$$\phi_2(x) = x^\eta (1 + |\log x|)^\beta$$

are regularly varying both at the origin (of indices $\min(\alpha, \beta)$ and η , respectively) and at infinity (of indices $\max(\alpha, \beta)$ and η , respectively).

Example 3.5. Functions ϕ are not confined to regularly varying functions. Indeed, $\phi(x) = x^\eta e^x$ satisfies $\text{WLSC}(\eta, 0, 1)$, but is not regularly varying at infinity.

3.2. Assumption $\text{DC}(\alpha, \gamma, d)$ on open sets. In what follows we assume that D is an open set in a metric measure space (X, ρ, μ) . We denote $\delta_x = \text{dist}(x, X \setminus D)$ for $x \in X$.

Definition 3.6. We say that D satisfies condition $\text{DC}(\alpha, \gamma, d)$, where $\gamma \in \mathbb{R}$, $d > 0$, $\alpha > 0$, $\alpha \neq 1$, if there exist $M > 0$ and (possibly empty) families $\mathcal{B}^{(n)} = \{B_j^{(n)}\}$ of subsets of D indexed by $n \in \mathbb{Z}$ such that the following conditions (B1)–(B4) hold.

(B1) $D = \cup_{j,n} B_j^{(n)}$ and each $x \in D$ belongs to at most M sets $B_j^{(n)}$.

(B2) For any $B_j^{(n)}$ we have

$$M^{-1}\alpha^n \leq \delta_x \leq M\alpha^n, \quad x \in B_j^{(n)},$$

$$M^{-1}\alpha^{nd} \leq \mu(B_j^{(n)}) \leq M\alpha^{nd}.$$

(B3) For any $B_j^{(n)}$ and any integer $k > M$, there exists a nonempty finite set $V(B_j^{(n)}, k)$ of indices so that, for each $i \in V(B_j^{(n)}, k)$,

$$\sup\{\rho(x, y) : x \in B_j^{(n)} \text{ and } y \in B_i^{(n+k)}\} \leq M(\alpha^n \vee \alpha^{n+k}).$$

(B4) For each $n \in \mathbb{Z}$ and $k > M$,

$$\sup_i \sum_{j:i \in V(B_j^{(n)}, k)} \frac{1}{\#V(B_j^{(n)}, k)} \leq M\alpha^{k\gamma}.$$

This definition is technical, but it allows to prove fractional Hardy inequality in different cases. Below we provide some illustrative examples of a set satisfying condition $\text{DC}(\alpha, \gamma, d)$. In the two examples $X = \mathbb{R}^d$ with the Euclidean distance, in which case $\delta_x = \text{dist}(x, \partial D)$ for all $x \in D$. Moreover, μ is the Lebesgue measure.

Example 3.7. Set $D = \mathbb{R}^d \setminus \{0\}$ satisfies condition $\text{DC}(\alpha, \gamma, d)$ with $\alpha = 2$, $\gamma = 0$ and

$$M = 2 \vee (1 - 2^{-d})|B(0, 1)| \vee \frac{1}{(1 - 2^{-d})|B(0, 1)|}.$$

Indeed, one may take $B_1^{(n)} := B(0, 2^n) \setminus B(0, 2^{n-1})$. That is, for each n there is exactly one set $B_j^{(n)}$, namely one with $j = 1$. Then $V(B_1^{(n)}, k) = \{1\}$ in (B3).

Example 3.8. Set $D = \mathbb{R}^d \setminus \{0\}$ satisfies condition $\text{DC}(\alpha, \gamma, d)$ with $\alpha = \frac{1}{2}$, $\gamma = 0$ and

$$M = 2 \vee (1 - 2^{-d})|B(0, 1)| \vee \frac{1}{(1 - 2^{-d})|B(0, 1)|}.$$

Indeed, one may take $B_1^{(n)} := B(0, 2^{-n}) \setminus B(0, 2^{-(n+1)})$ and $V(B_1^{(n)}, k) = \{1\}$ in (B3).

3.3. **Proof of Theorem 4.** Let us write

$$q = 2^{p+1} M^{4+2|\eta|} H^{-1} a^{k(\eta+\gamma-d)}, \quad R = 1 + M^2(1 \vee a^k), \quad S = 2^{p+1} a^{-kd} M^{d+1},$$

where $k > M$ is chosen such that $q < 1$ and $a^k \vee a^{-k} > M^2$.

We fix a function u for which the left hand side of (3.1) is finite, and define a set

$$F = \left\{ x \in D : |u(x)|^p > S \delta_x^{-d} \int_{D \cap B(x, R\delta_x)} |u(x) - u(y)|^p \mu(dy) \right\}.$$

Let us first observe that, for $x \in D \setminus F$,

$$(3.3) \quad \frac{|u(x)|^p}{\phi(\delta_x)} \leq S \int_{D \cap B(x, R\delta_x)} \frac{|u(x) - u(y)|^p}{\phi(\delta_x) \delta_x^d} \mu(dy).$$

Note that if the set F were empty, we would be already done.

At this stage we fix n and claim that, for $x \in F \cap B_j^{(n)}$ and $i \in V(B_j^{(n)}, k)$, we have

$$(3.4) \quad \mu \left(\left\{ y \in B_i^{(n+k)} : \frac{1}{2}|u(x)| \leq |u(y)| \leq \frac{3}{2}|u(x)| \right\} \right) \geq \frac{1}{2} \mu(B_i^{(n+k)}).$$

Suppose (3.4) fails. By our choice of R and conditions (B2) and (B3), $B_i^{(n+k)} \subset D \cap B(x, R\delta_x)$. Thus, we have

$$\begin{aligned} \int_{D \cap B(x, R\delta_x)} |u(x) - u(y)|^p \mu(dy) &\geq \int_{B_i^{(n+k)}} |u(x) - u(y)|^p \mu(dy) \\ &\geq \frac{1}{2} \mu(B_i^{(n+k)}) \cdot 2^{-p} |u(x)|^p \\ &\geq 2^{-p-1} a^{kd} M^{-d-1} \delta_x^d |u(x)|^p \\ &= S^{-1} \delta_x^d |u(x)|^p, \end{aligned}$$

which contradicts $x \in F$. Thus inequality (3.4) holds as claimed.

Let us record the following estimates for $B_i^{(n+k)} \in \mathcal{B}^{(n+k)}$ and $B_j^{(n)} \in \mathcal{B}^{(n)}$. By condition (B2), $\mu(B_j^{(n)}) \leq M^2 a^{-kd} \mu(B_i^{(n+k)})$, moreover, for $x \in B_j^{(n)}$ and $y \in B_i^{(n+k)}$ it holds $M^2 a^{-k} \delta_y \geq \delta_x \geq M^{-2} a^{-k} \delta_y$. Hence, by condition (3.2)

$$\phi(\delta_x) = \phi \left(\delta_y \frac{\delta_x}{\delta_y} \right) \geq H \left(\frac{\delta_x}{\delta_y} \right)^\eta \phi(\delta_y) \geq HM^{-2|\eta|} a^{-k\eta} \phi(\delta_y).$$

Here we need to ensure that $\frac{\delta_x}{\delta_y} < 1$ in the case when $a > 1$ and that $\frac{\delta_x}{\delta_y} > 1$ in the case when $a < 1$. But these are satisfied since, by assumption, $a^k \vee a^{-k} > M^2$, i.e., k is large enough. By the above estimate and inequality (3.4) we obtain

$$\int_{F \cap B_j^{(n)}} \frac{|u(x)|^p}{\phi(\delta_x)} \mu(dx) \leq \mu(B_j^{(n)}) \sup_{x \in F \cap B_j^{(n)}} \frac{|u(x)|^p}{\phi(\delta_x)}$$

$$\leq \frac{2^{p+1}M^2\alpha^{-kd}}{\#\mathcal{V}(B_j^{(n)}, k)} \sum_{i \in \mathcal{V}(B_j^{(n)}, k)} \int_{B_i^{(n+k)}} \frac{|u(y)|^p}{HM^{-2|n|}\alpha^{-kn}\phi(\delta_y)} \mu(dy).$$

After summing over all j

$$\begin{aligned} \sum_j \int_{F \cap B_j^{(n)}} \frac{|u(x)|^p}{\phi(\delta_x)} \mu(dx) &\leq 2^{p+1}M^{2+2|n|}H^{-1}\alpha^{k(n-d)} \times \sup_i \sum_{j: i \in \mathcal{V}(B_j^{(n)}, k)} \frac{1}{\#\mathcal{V}(B_j^{(n)}, k)} \\ &\times \sum_i \int_{B_i^{(n+k)}} \frac{|u(y)|^p}{\phi(\delta_y)} \mu(dy) \\ &\leq 2^{p+1}M^{3+2|n|}H^{-1}\alpha^{k(n+\gamma-d)} \sum_i \int_{B_i^{(n+k)}} \frac{|u(y)|^p}{\phi(\delta_y)} \mu(dy), \end{aligned}$$

and after summing over all n

$$\int_F \frac{|u(x)|^p}{\phi(\delta_x)} \mu(dx) \leq q \int_D \frac{|u(y)|^p}{\phi(\delta_y)} \mu(dy).$$

Recall that $q < 1$. Hence, by finiteness of the left hand side of (3.1),

$$\int_F \frac{|u(x)|^p}{\phi(\delta_x)} \mu(dx) \leq \frac{q}{1-q} \int_{D \setminus F} \frac{|u(y)|^p}{\phi(\delta_y)} \mu(dy).$$

This estimate and inequality (3.3) finish the proof. \square

3.4. Proof of Theorem 1. We use Potter's theorem [3, Theorem 1.5.6] to replace $\phi(\delta_x)$ by $c\phi(|x-y|)$ in the denominator, with $c = c(R, \phi)$. The assumption $0 < \rho_0, \rho_\infty$ is used here. The result follows now from Theorem 4 and examples 3.4, 3.7 and 3.8. \square

4. FAT BOUNDARY

During the course of this section, we prove Proposition 5. Then, in §4.4 we study the relations between lower Assouad dimension and uniform fatness.

Proposition 5. *Suppose $D \neq \emptyset$ is a proper κ -plump open set in \mathbb{R}^d such that, if D is unbounded, then ∂D is unbounded. Then D satisfies $DC(\alpha, \lambda, d)$ if $\alpha = 1/2$ and either $0 < \lambda < \underline{\dim}_A(\partial D)$ or $\lambda = 0$. Moreover, the associated constant M depends only on d, κ, λ and the constant C appearing in (F1) below.*

Under the assumptions of Proposition 5, the following two conditions (F1) and (F2) hold.

- (F1) There is a constant $C > 0$ as follows. Let $0 < r < R < 2\text{diam}(\partial D)$ and $x \in \partial D$. Suppose that B_1, \dots, B_N is a cover of $B(x, R) \cap \partial D$ by balls $B_j = B(\omega_j, r)$ with $\omega_j \in \partial D$ for $j = 1, \dots, N$. Then $N \geq C(R/r)^\lambda$.
- (F2) for each $0 < r < \text{diam}(D)$ and each $x \in \partial D$, there is $z \in \bar{B}(x, r)$ so that $B(z, \kappa r) \subset D$.

4.1. **Construction of families $\mathcal{B}^{(n)}$.** We define a constant

$$(4.1) \quad \tau = \left(\frac{15\sqrt{d}}{\kappa} \right)^d > 1.$$

For a given $n \in \mathbb{Z}$ and $\alpha \in \{\frac{1}{2}, 2\}$, we define

$$\mathcal{B}^{(n)} := \mathcal{B}_{1/2}^{(n)}, \quad \mathcal{B}_\alpha^{(n)} := \{\mathcal{B}_j^{(n)}\} := \{Q \in \mathcal{W}(D) : \tau^{-1} \leq \alpha^{-nd}|Q| \leq \tau\}.$$

Recall that $\mathcal{W}(D)$ stands for a Whitney decomposition of D . In particular, by inequalities (2.1), for any $x \in \mathcal{B}_j^{(n)} \in \mathcal{B}_\alpha^{(n)}$,

$$\tau^{-1/d}\alpha^n \leq \delta_x = \text{dist}(x, \partial D) \leq 5\sqrt{d}\tau^{1/d}\alpha^n.$$

Observe also that a given Whitney cube $Q \in \mathcal{W}(D)$ may belong to at most $1 + 2d^{-1}\log_2 \tau$ families $\mathcal{B}^{(n)} = \mathcal{B}_\alpha^{(n)}$ indexed by $n \in \mathbb{Z}$. Let us denote by $x_j^{(n)}$ the midpoint of $\mathcal{B}_j^{(n)}$. For later purposes we fix, once and for all, any point $y_j^{(n)} \in \partial D$ for which

$$|x_j^{(n)} - y_j^{(n)}| = \text{dist}(x_j^{(n)}, \partial D).$$

4.2. **Families $V(\mathcal{B}_j^{(n)}, k)$ for k large.** If D is unbounded, we construct families $V(\mathcal{B}_j^{(n)}, k)$ for $k > 3$. If D is bounded, then we construct these families for $k > 3 \vee \log_2(5\tau^{1/d})$.

Let us fix $\mathcal{B}_j^{(n)} \in \mathcal{B}^{(n)}$, and define $E := B(y_j^{(n)}, 2^{-n}) \cap \partial D$. By the $5r$ -covering theorem, see for instance [31, p. 23], there are points $\omega_1, \dots, \omega_N \in E$ such that the balls $B_m := B(\omega_m, 2^{-n-k})$ are disjoint and E is covered by the union of balls $5B_m$, $m = 1, \dots, N$. Let us estimate the number $N =: N_j^{(n,k)}$ of these balls;

Lemma 6. *We have $N_j^{(n,k)} \geq C5^{-\lambda}\tau^{-\lambda/d}2^{k\lambda}$.*

Proof. First consider the case when D is unbounded. Since $k > 3$, we find that

$$r := 5 \cdot 2^{-n-k} < 2^{-n} =: R.$$

Recall that the balls $5B_m = B(\omega_m, r)$ cover the set $E = B(y_j^{(n)}, R) \cap \partial D$. By condition (F1), we find that $N \geq C(R/r)^\lambda = C5^{-\lambda}2^{k\lambda}$. The bounded case is similar, and we use the facts that $k > 3 \vee \log_2(5\tau^{1/d})$ and $\text{diam}(\partial D) \geq \text{diam}(D)$. \square

The next step is to use the plumpness condition (F2) in order to locate a sufficiently large cube inside each B_m . Namely, for each $m = 1, \dots, N$, there is $z_m \in \bar{B}(\omega_m, 2^{-n-k}/3)$ such that

$$B(z_m, \kappa 2^{-n-k}/3) \subset D.$$

Let us consider a Whitney cube $Q_m \in \mathcal{W}(D)$ for which $z_m \in Q_m$. By inequalities (2.1), we have $Q_m \subset B_m$. Moreover,

$$\kappa 2^{-n-k}/3 \leq \text{dist}(z_m, \partial D) \leq 5\text{diam}(Q_m) \leq 5\text{dist}(z_m, \partial D) \leq 5 \cdot 2^{-n-k}/3.$$

Hence, by our definition (4.1) of τ , we obtain

$$\tau^{-1} \leq 2^{d(n+k)}|Q_m| \leq \tau.$$

That is, cube $Q_m \subset B_m$ belongs to $\mathcal{B}^{(n+k)}$. Since the balls B_m , $m = 1, \dots, N$, are disjoint, also the cubes Q_m are disjoint. Hence, the indexing set

$$V(B_j^{(n)}, k) = \{i : B_i^{(n+k)} = Q_m \text{ for some } m = 1, \dots, N_j^{(n,k)}\}$$

contains exactly $N_j^{(n,k)}$ indices.

4.3. Proof of Proposition 5. We focus on conditions (B3) and (B4), as the remaining conditions are clearly satisfied. Let us fix $B_j^{(n)}$ and k large enough so that $V(B_j^{(n)}, k)$ is defined. Let us consider $i \in V(B_j^{(n)}, k)$, and two given points $x \in B_j^{(n)}$ and $y \in B_i^{(n+k)}$. Using the notation above, we have $B_i^{(n+k)} = Q_m \subset B_m$ for some $m = 1, \dots, N_j^{(n,k)}$. Thus,

$$\begin{aligned} |x - y| &\leq |x - x_j^{(n)}| + |x_j^{(n)} - y_j^{(n)}| + |y_j^{(n)} - \omega_m| + |\omega_m - y| \\ &< \text{diam}(B_j^{(n)}) + \text{dist}(x_j^{(n)}, \partial D) + 2^{-n} + 2^{-n-k} \\ &\leq 8\sqrt{d}\tau^{1/d}2^{-n}. \end{aligned}$$

This is condition (B3). A particular consequence of this estimate is the following. We fix a cube $B_i^{(n+k)}$ and a point y therein. Then, if $B_j^{(n)} \in \mathcal{B}^{(n)}$ is such that $i \in V(B_j^{(n)}, k)$,

$$B_j^{(n)} \subset B(y, 8\sqrt{d}\tau^{1/d}2^{-n}).$$

Since the interiors of cubes in $\mathcal{B}^{(n)}$ are disjoint, we find that there are at most

$$\frac{(16\sqrt{d}\tau^{1/d}2^{-n})^d}{\tau^{-1}2^{-nd}} = (16\sqrt{d})^d \tau^2$$

cubes $B_j^{(n)}$ subject to the conditions above. By using this fact, we may now deduce the remaining estimate as follows; For a fixed i ,

$$\sum_{j:i \in V(B_j^{(n)}, k)} \frac{1}{\#V(B_j^{(n)}, k)} = \sum_{j:i \in V(B_j^{(n)}, k)} \frac{1}{N_j^{(n,k)}} \leq (16\sqrt{d})^d \tau^{2+\lambda/d} C^{-1} 5^\lambda 2^{-k\lambda}.$$

This is condition (B4). □

4.4. Lower Assouad dimension and uniform fatness. We provide a useful connection between the lower Aikawa dimension and (local) uniform fatness. For further discussion, we refer to [19]. Uniform fatness is usually defined in terms of Riesz capacities, [1, 26]. In case of closed sets, there is an equivalent definition—in terms of Hausdorff content—that we adopt. This equivalence is based on the self-improving properties of closed uniformly fat sets, [16].

Recall that the λ -Hausdorff content of a set $E \subset \mathbb{R}^d$ is

$$\mathcal{H}_\infty^\lambda(E) = \inf \left\{ \sum_{i=1}^{\infty} r_i^\lambda : E \subset \bigcup_{i=1}^{\infty} B(x_i, r_i), r_i > 0 \right\}.$$

As is easily seen, we may allow also finite coverings in the infimum above. Let $1 < p < \infty$ and $0 < s < d/p$. We say that the boundary ∂D is (s, p) -uniformly fat, if there is $d - sp < \lambda \leq d$ and a constant $C > 0$ such that

$$(4.2) \quad \mathcal{H}_\infty^\lambda(B(x, R) \cap \partial D) \geq CR^\lambda$$

for all $x \in \partial D$ and $R > 0$.

Note that ∂D and D have to be unbounded if the boundary is (s, p) -uniformly fat. Remark 2.3 in [19] shows that $\underline{\dim}_A(\partial D)$ is the supremum of all $\lambda \geq 0$ for which (4.2) holds for every $x \in \partial D$ and $0 < R < \text{diam}(\partial D)$. Below, for the convenience of the reader, we provide a detailed treatment of certain consequences of this statement—that are needed for Corollary 3.

Proposition 7. *Suppose D is an open set in \mathbb{R}^d so that ∂D is (s, p) -uniformly fat for $1 < p < \infty$ and $0 < s < d/p$. Then D satisfies condition (F1) for some $d - sp < \lambda \leq d$ and, as a consequence, we have a strict inequality $\underline{\dim}_A(\partial D) > d - sp$.*

Proof. By assumption, there is $d - sp < \lambda \leq d$ and $C > 0$ such that (4.2) holds for all $x \in \partial D$ and $R > 0$. Let us fix $x \in \partial D$ and $0 < r < R$. Suppose that B_1, \dots, B_N is a cover of $B(x, R) \cap \partial D$ by balls $B_j = B(\omega_j, r)$ with $\omega_j \in \partial D$. Then, by (4.2),

$$Nr^\lambda = \sum_{j=1}^N r^\lambda \geq \mathcal{H}_\infty^\lambda(B(x, R) \cap \partial D) \geq CR^\lambda.$$

Thus, $N \geq C(R/r)^\lambda$, as required. \square

As we have observed, the uniform fatness is a convenient notion in case of unbounded open sets. In case of a bounded open set D in \mathbb{R}^d , it is natural to assume that ∂D is (s, p) -locally uniformly fat. That is, there is $d - sp < \lambda \leq d$ and a constant $C > 0$ such that inequality (4.2) holds for all $x \in \partial D$ and $0 < R < 2 \text{diam}(\partial D) < \infty$.

The following result is analogous to Proposition 7.

Proposition 8. *Let D be a bounded open set in \mathbb{R}^d such that ∂D is (s, p) -locally uniformly fat for $1 < p < \infty$ and $0 < s < d/p$. Then D satisfies condition (F1) for some $d - sp < \lambda \leq d$ and, as a consequence, we have a strict inequality $\underline{\dim}_A(\partial D) > d - sp$.*

Example 4.1. Consider the Koch snowflake domain $D \subset \mathbb{R}^2$. It is a bounded κ -plump domain and ∂D is (s, p) -locally uniformly fat if $1 < p < \infty$ and $2 - \log 4 / \log 3 < sp < 2$, see e.g. [16].

5. THIN BOUNDARY

The main result in this section is the following.

Proposition 9. *Let $D \neq \emptyset$ be an unbounded κ -plump open set in \mathbb{R}^d , $D \neq \mathbb{R}^d$. Then D satisfies condition DC(α, λ, d) for $\alpha = 2$ and $\lambda > \overline{\dim}_A(\partial D)$. The associated constant M depends only on d, κ, λ and the constant C appearing in (T1) below.*

Before the proof, let us clarify the assumptions. Under the assumptions of Proposition 9, the following two conditions (T1) and (T2) hold.

(T1) there is a constant $C > 0$ as follows. Assuming that $0 < r < R$ and $x \in \partial D$, there is a cover of $B(x, R) \cap \partial D$ by using balls $B(\omega_j, r)$ with $\omega_j \in \partial D$, $j = 1, \dots, N$, such that the number of these balls satisfies inequality $N \leq C(R/r)^\lambda$.

(T2) for each $0 < r$ and each $x \in \partial D$, there is a point $z \in \bar{B}(x, r)$ such that $B(z, \kappa r) \subset D$.

We note that in Definition 2.2 the restriction $R < 2 \operatorname{diam}(E)$ may be removed, resulting in no such restriction in (T1).

5.1. **Construction of families $\mathcal{B}^{(n)}$.** Let $\tau > 1$ be defined by (4.1). For $n \in \mathbb{Z}$, we define

$$\mathcal{B}^{(n)} := \mathcal{B}_2^{(n)} := \{B_j^{(n)}\} := \{Q \in \mathcal{W}(D) : \tau^{-1} \leq 2^{-nd}|Q| \leq \tau\}.$$

For properties of cubes in these families and definition of $y_j^{(n)}$, see §4.1. Recall also that $\mathcal{W}(D)$ is a Whitney decomposition of D , we refer to §2.

5.2. **Families $V(B_j^{(n)}, k)$ with $k > 0$.** Let us fix a cube $B_j^{(n)} \in \mathcal{B}^{(n)}$. By condition (T2), there is a point $z_j^{(n)} \in \bar{B}(y_j^{(n)}, 2^{n+k})$ such that

$$B(z_j^{(n)}, \kappa 2^{n+k}) \subset D.$$

Observe how the unboundedness of D is visible here, as $k > 0$ is arbitrary. Let $Q_j \in \mathcal{W}(D)$ be a Whitney cube such that $z_j^{(n)} \in Q_j$. Then

$$\kappa \cdot 2^{n+k} \leq \operatorname{dist}(z_j^{(n)}, \partial D) \leq 5 \operatorname{diam}(Q_j) \leq 5 \operatorname{dist}(z_j^{(n)}, \partial D) \leq 5 \cdot 2^{n+k}.$$

By definition of (4.1) of τ , we then have $Q_j = B_i^{(n+k)} \in \mathcal{B}^{(n+k)}$ for some index i . We define

$$V(B_j^{(n)}, k) = \{i\}.$$

5.3. **Proof of Proposition 9.** We need an auxiliary estimate analogous to [19, Lemma 4.3]; condition (T1) is our primary tool. For $m \in \mathbb{Z}$, $\omega \in \partial D$, and $R > 0$, we denote

$$\mathcal{W}_m(D; B(\omega, R)) = \{Q \in \mathcal{W}_m(D) : Q \subset B(\omega, R)\}.$$

Lemma 10. *Let $0 < 2^{-m} \leq R$, where $m \in \mathbb{Z}$. Then for every $\omega \in \partial D$,*

$$\#\mathcal{W}_m(D; B(\omega, R)) \leq C(14\sqrt{d})^{d+\lambda} \left(\frac{R}{2^{-m}}\right)^\lambda.$$

Proof. Suppose B_1, \dots, B_N is a cover of $B(\omega, 6\sqrt{d}R) \cap \partial D$ by balls $B_j = B(\omega_j, 2^{-m})$ that are centred in ∂D , see condition (T1). Consider a cube $Q \in \mathcal{W}_m(D; B(\omega, R))$, and fix a point $y_Q \in \partial D$ such that $|x_Q - y_Q| = \operatorname{dist}(x_Q, \partial D)$. Here x_Q denotes the midpoint of Q . By inequalities (2.1) and the fact that $Q \subset B(\omega, R)$,

$$|y_Q - \omega| \leq |y_Q - x_Q| + |x_Q - \omega| < 5 \operatorname{diam}(Q) + R \leq 6\sqrt{d}R.$$

By the covering property, there is $j = j(Q)$ such that $y_Q \in B_j$. We can infer that

$$\mathcal{W}_m(D; B(\omega, R)) = \bigcup_{j=1}^N Q_j,$$

where $\mathcal{Q}_j = \{Q \in \mathcal{W}_m(D; B(\omega, R)) : y_Q \in B_j\}$. Let $Q \in \mathcal{Q}_j$. Then, for every $x \in Q$,

$$\begin{aligned} |x - \omega_j| &\leq |x - x_Q| + |x_Q - y_Q| + |y_Q - \omega_j| \\ &< \text{diam}(Q) + \text{dist}(x_Q, \partial D) + 2^{-m} \\ &\leq 7\sqrt{d}2^{-m}. \end{aligned}$$

Since the interiors of cubes in the family \mathcal{Q}_j are disjoint, there are at most

$$\frac{|B(\omega_j, 7\sqrt{d}2^{-m})|}{2^{-md}} \leq (14\sqrt{d})^d$$

cubes in this family. Hence,

$$\#\mathcal{W}_m(D; B(\omega, R)) \leq \sum_{j=1}^N \#\mathcal{Q}_j \leq (14\sqrt{d})^d N \leq C \left(\frac{6\sqrt{d}R}{2^{-m}} \right)^\lambda \cdot (14\sqrt{d})^d.$$

This concludes the proof. \square

We are ready to prove the main result in this section.

Proof of Proposition 9. The properties (B1) and (B2) are clear. In order to verify condition (B3), let us fix $k > 0$ and a cube $B_j^{(n)}$. Consider $i \in V(B_j^{(n)}, k)$, and points $x \in B_j^{(n)}$ and $z \in B_i^{(n+k)}$. Then, by the construction above,

$$\begin{aligned} |z - x| &\leq |z - z_j^{(n)}| + |z_j^{(n)} - y_j^{(n)}| + |y_j^{(n)} - x_j^{(n)}| + |x_j^{(n)} - x| \\ &\leq \text{diam}(B_i^{(n+k)}) + 2^{n+k} + \text{dist}(x_j^{(n)}, \partial D) + \text{diam}(B_j^{(n)}) \\ &< 8\sqrt{d}\tau^{1/d}2^{n+k}. \end{aligned}$$

This is condition (B3).

In order to verify the last condition (B4), we fix cubes $B_i^{(n+k)}$ and $B_j^{(n)}$ such that $i \in V(B_j^{(n)}, k)$. Then $\#\mathcal{V}(B_j^{(n)}, k) = 1$. Moreover,

$$B_j^{(n)} \subset B(y_i^{(n+k)}, 13\sqrt{d}\tau^{1/d}2^{n+k}).$$

Indeed, for any $x \in B_j^{(n)}$,

$$|x - y_i^{(n+k)}| \leq |x - x_i^{(n+k)}| + |x_i^{(n+k)} - y_i^{(n+k)}| < 13\sqrt{d}\tau^{1/d}2^{n+k}.$$

We still need another auxiliary estimate, namely, if $m \in \mathbb{Z}$ is such that $B_j^{(n)} \in \mathcal{W}_m(D)$, then $\tau^{-1/d} \leq 2^{m+n} \leq \tau^{1/d}$. We can finally proceed as follows

$$\begin{aligned} \sum_{j:i \in V(B_j^{(n)}, k)} \frac{1}{\#\mathcal{V}(B_j^{(n)}, k)} &= \#\{j : i \in V(B_j^{(n)}, k)\} \\ &= \sum_m \#\{j : i \in V(B_j^{(n)}, k) \text{ and } B_j^{(n)} \in \mathcal{W}_m(D)\} \end{aligned}$$

$$\begin{aligned} &\leq \sum_m \#\mathcal{W}_m(D; B(y_i^{(n+k)}, 13\sqrt{d}\tau^{1/d}2^{n+k})) \\ &\leq \sum_m C(14\sqrt{d})^{d+\lambda} \left(\frac{13\sqrt{d}\tau^{1/d}2^{n+k}}{2^{-m}} \right)^\lambda, \end{aligned}$$

where m ranges over indices $-n - \log_2 \tau^{1/d} \leq m \leq -n + \log_2 \tau^{1/d}$. This yields condition (B4). \square

REFERENCES

- [1] D. R. Adams and L. I. Hedberg. *Function spaces and potential theory*, volume 314 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1996.
- [2] P. Assouad. Plongements lipschitziens dans \mathbf{R}^n . *Bull. Soc. Math. France*, 111(4):429–448, 1983.
- [3] N. H. Bingham, C. M. Goldie, and J. L. Teugels. *Regular variation*, volume 27 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1989.
- [4] K. Bogdan and B. Dyda. The best constant in a fractional Hardy inequality. *Math. Nachr.*, 284(5-6):629–638, 2011.
- [5] K. Bogdan, T. Grzywny, and M. Ryznar. Density and tails of unimodal convolution semigroups. arXiv:1305.0976 [math.FA], 2013.
- [6] G. Bouligand. Sur quelques points de topologie restreinte du premier ordre. *Bull. Soc. Math. France*, 56:26–35, 1928.
- [7] B. Dyda. A fractional order Hardy inequality. *Illinois J. Math.*, 48(2):575–588, 2004.
- [8] B. Dyda. Fractional Hardy inequality with a remainder term. *Colloq. Math.*, 122(1):59–67, 2011.
- [9] B. Dyda and R. L. Frank. Fractional Hardy-Sobolev-Maz'ya inequality for domains. *Studia Math.*, 208(2):151–166, 2012.
- [10] D. Edmunds, R. Hurri-Syrjänen, and A. V. Vähäkangas. Fractional Hardy-type inequalities in domains with uniformly fat complement. *Proc. Amer. Math. Soc.*, to appear.
- [11] S. Filippas, L. Moschini, and A. Tertikas. Sharp Trace Hardy–Sobolev-Maz'ya Inequalities and the Fractional Laplacian. *Arch. Ration. Mech. Anal.*, 208(1):109–161, 2013.
- [12] R. L. Frank and R. Seiringer. Sharp fractional Hardy inequalities in half-spaces. In *Around the research of Vladimir Maz'ya. I*, volume 11 of *Int. Math. Ser. (N. Y.)*, pages 161–167. Springer, New York, 2010.
- [13] T. Grzywny. On Harnack inequality and Hölder regularity for isotropic unimodal Lévy processes. arXiv:1301.2441 [math.PR], 2013.
- [14] P. Gurka and B. Opic. Sharp Hardy inequalities of fractional order involving slowly varying functions. *J. Math. Anal. Appl.*, 386(2):728–737, 2012.
- [15] H. P. Heinig, A. Kufner, and L.-E. Persson. On some fractional order Hardy inequalities. *J. Inequal. Appl.*, 1(1):25–46, 1997.
- [16] L. Ichnatsyeva, J. Lehrbäck, H. Tuominen, and A. V. Vähäkangas. Fractional Hardy inequalities and visibility of the boundary. arXiv:1305.4616.
- [17] L. Ichnatsyeva and A. V. Vähäkangas. Hardy inequalities in Triebel-Lizorkin spaces. *Indiana Univ. Math. J.*, to appear.
- [18] L. Ichnatsyeva and A. V. Vähäkangas. Hardy inequalities in Triebel-Lizorkin spaces II. Aikawa dimension. arXiv:1302.5295.
- [19] A. Käenmäki, J. Lehrbäck, and M. Vuorinen. Dimensions, Whitney covers, and tubular neighborhoods. *Indiana Univ. Math. J.*, to appear.
- [20] P. Kim and A. Mimica. Harnack inequalities for subordinate Brownian motions. *Electron. J. Probab.*, 17:no. 37, 23, 2012.

- [21] P. Koskela and X. Zhong. Hardy's inequality and the boundary size. *Proc. Amer. Math. Soc.*, 131(4):1151–1158 (electronic), 2003.
- [22] A. Kufner and L.-E. Persson. *Weighted inequalities of Hardy type*. World Scientific Publishing Co. Inc., River Edge, NJ, 2003.
- [23] J. Lehrbäck. Hardy inequalities and Assouad dimensions. In preparation.
- [24] J. Lehrbäck. Weighted Hardy inequalities beyond Lipschitz domains. *Proc. Amer. Math. Soc.*, to appear, arXiv:1209.0588.
- [25] J. Lehrbäck and H. Tuominen. A note on the dimensions of Assouad and Aikawa. *J. Math. Soc. Japan*, 65(2):343–356, 2013.
- [26] J. L. Lewis. Uniformly fat sets. *Trans. Amer. Math. Soc.*, 308(1):177–196, 1988.
- [27] M. Loss and C. Sloane. Hardy inequalities for fractional integrals on general domains. *J. Funct. Anal.*, 259(6):1369–1379, 2010.
- [28] J. Luukkainen. Assouad dimension: antifractal metrization, porous sets, and homogeneous measures. *J. Korean Math. Soc.*, 35(1):23–76, 1998.
- [29] O. Martio. Definitions for uniform domains. *Ann. Acad. Sci. Fenn. Ser. A I Math.*, 5(1):197–205, 1980.
- [30] O. Martio and J. Väisälä. Bounded turning and passability. *Results Math.*, 24:246–273, 1993.
- [31] P. Mattila. *Geometry of sets and measures in Euclidean spaces*, volume 44 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1995. Fractals and rectifiability.
- [32] J. Nečas. Sur une méthode pour résoudre les équations aux dérivées partielles du type elliptique, voisine de la variationnelle. *Ann. Scuola Norm. Sup. Pisa (3)*, 16:305–326, 1962.
- [33] H. Šikić, R. Song, and Z. Vondraček. Potential theory of geometric stable processes. *Probab. Theory Related Fields*, 135(4):547–575, 2006.
- [34] C. A. Sloane. A fractional Hardy-Sobolev-Maz'ya inequality on the upper halfspace. *Proc. Amer. Math. Soc.*, 139(11):4003–4016, 2011.
- [35] E. M. Stein. *Singular integrals and differentiability properties of functions*. Princeton Mathematical Series, No. 30. Princeton University Press, Princeton, N.J., 1970.
- [36] J. Väisälä. Uniform domains. *Tohoku Math. J. (2)*, 40(1):101–118, 1988.
- [37] A. Wannebo. Hardy inequalities. *Proc. Amer. Math. Soc.*, 109(1):85–95, 1990.

(B.D.) FACULTY OF MATHEMATICS, UNIVERSITY OF BIELEFELD, POSTFACH 10 01 31, D-33501 BIELEFELD, GERMANY, AND INSTITUTE OF MATHEMATICS AND COMPUTER SCIENCE, WROCLAW UNIVERSITY OF TECHNOLOGY, WYBRZEŻE WYSPIAŃSKIEGO 27, 50-370 WROCLAW, POLAND
E-mail address: dyda@math.uni-bielefeld.de bdyda@pwr.wroc.pl

(A.V.V.) DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. Box 68, FI-00014 UNIVERSITY OF HELSINKI, FINLAND
E-mail address: antti.vahakangas@helsinki.fi